

Interface Yang-Mills, Supersymmetry, and Janus

Eric D'Hoker, John Estes and Michael Gutperle

*Department of Physics and Astronomy
University of California, Los Angeles, CA 90095, USA*

Abstract

We consider theories consisting of a planar interface with $\mathcal{N} = 4$ super-Yang-Mills on either side and varying gauge coupling across the interface. The interface does not carry any independent degrees of freedom, but is allowed to support local gauge invariant operators, included with independent interface couplings. In general, both conformal symmetry and supersymmetry will be broken, but for special arrangements of the interface couplings, these symmetries may be restored. We provide a systematic classification of all allowed interface supersymmetries. We find new theories preserving eight and four Poincaré supersymmetries, which get extended to sixteen and eight supersymmetries in the conformal limit, respectively with $SU(2) \times SU(2)$, $SO(2) \times SU(2)$ internal symmetry. The Lagrangians for these theories are explicitly constructed. We also recover the theory with two Poincaré supersymmetries and $SU(3)$ internal symmetry proposed earlier as a candidate CFT dual to super Janus. Since our new interface theories have only operators from the supergravity multiplet turned on, dual supergravity solutions are expected to exist. We speculate on the possible relation between the interface theory with maximal supersymmetry and the near-horizon limit of the D3-D5 system.

1 Introduction

The Janus solutions of Type IIB supergravity, found in [1], form a continuous family of deformations of $AdS_5 \times S_5$, in which the dilaton is allowed to vary along a single space direction, but the anti-symmetric 2-form field $B_{(2)}$ vanishes. The Janus solutions are smooth and singularity-free. They may be simply generalized to include a varying axion along with a varying dilaton, and admit an analytic representation in terms of elliptic functions [2]. They are invariant under $SO(2, 3) \times SO(6)$, but break all supersymmetries. Remarkably, despite their lack of supersymmetry, the solutions are classically stable against all small and a certain class of large perturbations [1, 3, 4].

The Janus solutions offer interesting candidates for further exploring the AdS/CFT correspondence [5, 6, 7] (for reviews, see [8, 9]). Some aspects of the holographic properties of the Janus solution were discussed in [1, 3, 10]. The boundary of a Janus solution consists of two halves of Minkowski space-time which are joined along an interface. The dilaton takes two different asymptotic values on each of these boundary components. The CFT dual, proposed in [1, 11], consists of a 3+1-dimensional gauge theory with a 2+1-dimensional planar interface. The gauge theory on each side of the planar interface is $\mathcal{N} = 4$ super Yang-Mills and the gauge coupling varies discontinuously across the interface. The two values of the gauge coupling correspond to the two asymptotic values of the dilaton in the Janus solution. The non-constancy of the gauge coupling allows the action to include *interface operators*, whose support is limited to the interface. These operators involve only $\mathcal{N} = 4$ super-Yang-Mills fields, and the interface carries no independent degrees of freedom. This is to be contrasted with defect or boundary conformal field theories [12, 13, 14], where often new degrees of freedom are localized on the defect.

At present, it is unclear whether the Janus solution arises as the near-horizon limit of any interesting brane configuration. The absence of independent degrees of freedom on the interface seems to preclude the presence of open strings between the D3 branes of the original undeformed $AdS_5 \times S_5$ and the interface. Or, if open strings did arise, the question becomes how they could have decoupled in the supergravity limit.

The absence of any degree of supersymmetry in the Janus solution makes it difficult to investigate such brane candidates directly. Therefore, one is interested in generalizing the original non-supersymmetric Janus to new solutions of Type IIB supergravity which do have non-trivial supersymmetry. Part of this goal was achieved in [11] where a supersymmetric Janus solution was found in a five dimensional gauged supergravity theory with $N = 2$ supersymmetry. In [2] supersymmetric Janus solutions of ten dimensional Type IIB supergravity were found with $\mathcal{N} = 1$ interface supersymmetry and $SU(3)$ in-

ternal symmetry. Clearly, however, it would be advantageous to understand not just one or a few such solutions, but rather the full space of supersymmetric Janus-like solutions. Progress in this direction can be made by analyzing the degree of supersymmetry in the dual CFT.

In this paper, we shall pose and solve the following problem. Consider a gauge theory in 3+1 dimensions with a 2+1-dimensional planar interface, with $\mathcal{N} = 4$ super Yang-Mills on each side of the interface, and no independent degrees of freedom on the interface. The action of the theory allows for local gauge invariant interface operators consistent with interface conformal invariance, to be included with arbitrary *interface couplings*. The problem is to classify completely, as a function of the interface couplings, the degree of supersymmetry and conformal invariance of the theory. Our investigation is close in spirit to [15, 16, 17].

A gauge coupling which is strictly discontinuous across the interface leads to a number of technical subtleties and complications, such as the possibility of discontinuous canonical fields and the appearance of the square of Dirac δ -functions [11]. Therefore, the problem posed in the preceding paragraph will be solved in two stages. In the first stage, the gauge coupling will be allowed to vary smoothly across the interface, and the problem of determining the degree of Poincaré supersymmetry in the presence of this smoothly varying gauge coupling will be solved. The supersymmetry transformation rules in the presence of the interface will require certain modifications from their standard form in $\mathcal{N} = 4$ super Yang-Mills. In the second stage, the existence of the limit of this smoothly varying gauge coupling to a discontinuous jump will be investigated. When this limit exists, interface conformal invariance will be recovered, and Poincaré supersymmetry will be enhanced to interface superconformal symmetry.

The results obtained in this paper may be summarized as follows. The conditions for Poincaré supersymmetry in the presence of a smoothly varying gauge coupling are reduced to a complicated looking set of 7 algebraic matrix equations involving the gauge coupling, the interface couplings, and the supersymmetry parameters. Remarkably, these equations may be drastically simplified and their solutions classified systematically. The existence of the theory [11] with 2 real Poincaré supercharges, $SU(3)$ global symmetry, and interface conformal invariance is confirmed. New solutions with 16 and 8 conformal supersymmetries are discovered, with respectively $SU(2) \times SU(2)$ and $SO(2) \times SU(2)$ symmetry.

The rest of the paper is organized as follows. In section 2, we define the general interface $\mathcal{N} = 4$ Yang-Mills theory, list its allowed interface terms and couplings, and

derive its global symmetries. In section 3, we derive a set of algebraic equations for the existence of at least one interface supersymmetry, which is solved in section 4. This result is used in section 5 to produce a systematic classification of allowed Poincaré interface supersymmetries. The existence of the conformal limit is discussed in section 6. In section 7, a comparison is made between the new theory with 16 interface supersymmetries and $SU(2) \times SU(2)$ internal symmetry, and the near-horizon limit of the D3-D5 system.

2 $\mathcal{N} = 4$ super Yang-Mills with an interface

In this section, we obtain a systematic generalization of the interface Yang-Mills theories, proposed in [1, 11] as duals to the Janus solution. The goal of the generalization is to achieve a complete classification of the possible supersymmetries supported by such interface theories. The generalization consists of 3+1-dimensional gauge theory, with a 2+1-dimensional planar interface, $\mathcal{N} = 4$ super Yang-Mills on each side of the interface, no independent degrees of freedom on the interface, and local gauge invariant interface operators included in the action with arbitrary interface couplings. Since we are primarily interested in conformal theories, we shall assume throughout that no dimensionful couplings enter into the action. This requirement also ensures renormalizability.

We start with a summary of standard $\mathcal{N} = 4$ super Yang-Mills [19, 20]. The theory contains a gauge field A_μ , four Weyl fermions ψ^a , and six real scalars ϕ^i , which transform under $SU(4)$ in the **1**, **4**, and **6** representations respectively, and under the $SU(N)$ gauge group in the adjoint representation.¹ The Lagrangian is given by

$$\begin{aligned} \mathbb{L}_0 = & -\frac{1}{4g^2} \text{tr}(F^{\mu\nu} F_{\mu\nu}) - \frac{1}{2g^2} \text{tr}(D^\mu \phi^i D_\mu \phi^i) + \frac{1}{4g^2} \text{tr}([\phi^i, \phi^j][\phi^i, \phi^j]) \\ & - \frac{i}{2g^2} \text{tr}(\bar{\psi} \gamma^\mu D_\mu \psi) + \frac{i}{2g^2} \text{tr}(D_\mu \bar{\psi} \gamma^\mu \psi) \\ & + \frac{1}{2g^2} \text{tr}(\psi^t \mathcal{C} \rho^i [\phi^i, \psi] + \psi^\dagger \mathcal{C} (\rho^i)^* [\phi^i, \psi^*]) \end{aligned} \quad (2.1)$$

The trace is over the gauge algebra only. The γ^μ are the Dirac matrices, and $\mathcal{C}$ is the associated charge conjugation matrix defined by $(\gamma^\mu)^t = -\mathcal{C} \gamma^\mu \mathcal{C}^{-1}$. The 4×4 matrices ρ^i are Clebsch-Gordan coefficients for the $SU(4)$ tensor product decomposition $\mathbf{4} \otimes \mathbf{4} \rightarrow \mathbf{6}$. Along with other useful $SU(4)$ matrices, the ρ^i are presented in Appendix A, where they are obtained from the Clifford algebra in 6 Euclidean dimensions. The gauge coupling g is a constant and the CP -violating $\text{tr}(F\tilde{F})$ term has been omitted.

The Lagrangian $\mathbb{L}_0$ is invariant under $SU(4)$ R-symmetry, conformal $SO(2, 4)$, and $\mathcal{N} = 4$ Poincaré supersymmetry, which is enhanced to $\mathcal{N} = 4$ conformal supersymmetry. These symmetries combine into a simple supergroup $PSU(2, 2|4)$.

A planar interface, whose position is described by the vanishing of a single space coordinate, x^π , may be introduced by making the coupling discontinuous across $x^\pi = 0$,

¹The fields A_μ , ψ^a and ϕ^i take values in the gauge algebra. Internal labels will be denoted by Latin indices $a = 1, \dots, 4$, and $i = 1, \dots, 6$. Each Weyl spinor ψ^a will be expressed as a 4-component Dirac spinor whose right chirality component vanishes, and the 4 Weyl spinors ψ^a will be grouped in a quadruplet for which we use the matrix notation ψ .

and adding local gauge invariant interface operators of dimension 3 with arbitrary interface couplings, but no independent degrees of freedom on the interface. In the context of the AdS/CFT correspondence, such an interface will not allow strings stretching between the interface and D3-branes. With this set-up, the interface will preserve gauge invariance, as well as the $SO(2,3)$ conformal invariance which leaves the plane $x^\pi = 0$ invariant. For general interface couplings, $SU(4)$ and supersymmetry will be broken.

It was remarked already in [11], however, that a strictly discontinuous gauge coupling may introduce technical complications, such as discontinuous canonical fields and squares of Dirac δ -functions at $x^\pi = 0$. Therefore, following [11], we introduce a gauge coupling function $g(x^\pi)$ which varies smoothly across the interface, such as represented schematically in Fig 1. In the presence of a smooth gauge coupling, no technical subtleties will arise, and the interface theory will be well-defined. It will be natural, not to localize interface operators strictly at $x^\pi = 0$, but instead to introduce them as 4-dimensional operators whose coupling is proportional to the derivative of the gauge coupling $\partial_\pi g$. Of course, any smooth, non-constant, $g(x^\pi)$ has at least one length scale and therefore will break $SO(2,3)$ conformal invariance further to 2+1-dimensional Poincaré invariance. Conformal invariance may then be recovered if the limit in which $g(x^\pi)$ tends to a step function actually exists and makes sense. It is interesting to note that our supersymmetry analysis will hold for a general gauge coupling varying along x^π .




Figure 1: The discontinuous gauge coupling is replaced by a smoothly varying function.

It remains to give a complete description of the local gauge invariant interface operators and their couplings. We introduce a length scale only through the gauge coupling function $g(x^\pi)$, so that the interface operators will always be multiplied by at least one

power of $\partial_\pi g$. This will guarantee that the interface Lagrangian will not contribute to the dynamics in the bulk. We write the full Lagrangian as follows,

$$\mathbb{L} = \mathbb{L}_0 + \mathbb{L}_I \quad (2.2)$$

where $\mathbb{L}_I$ stands for the interface Lagrangian. We now list the possible interface operators in $\mathbb{L}_I$, consistent with gauge invariance, 2+1-dimensional Poincaré invariance, and of dimension no higher than 3. It is useful to organize the contributions as follows,

$$\begin{aligned} \mathbb{L}_\psi &= \frac{(\partial_\pi g)}{g^3} \text{tr} \left(y_1 \bar{\psi} \gamma^\pi \psi + \frac{i}{4} y_2^{ij} \bar{\psi} \gamma^\pi \rho^{ij} \psi - \frac{i}{2} y_3^{ijk} \left(\psi^t \mathcal{C} \rho^{ijk} \psi + \psi^\dagger \mathcal{C} (\rho^{ijk})^* \psi^* \right) \right) \\ \mathbb{L}_\phi &= \frac{(\partial_\pi g)}{2g^3} \text{tr} \left(z_1^{ij} \partial_\pi (\phi^i \phi^j) + 2z_2^{ij} \phi^{[i} D_\pi \phi^{j]} - i z_3^{ijk} \phi^i [\phi^j, \phi^k] \right) \\ \mathbb{L}_{\phi^2} &= \frac{(\partial_\pi g)^2}{2g^4} z_4^{ij} \text{tr} (\phi^i \phi^j) \\ \mathbb{L}_{\text{CS}} &= \frac{(\partial_\pi g)}{g^3} z_{CS} \varepsilon^{\pi\mu\nu\rho} \text{tr} \left(A_\mu \partial_\nu A_\rho + \frac{2}{3} i A_\mu A_\nu A_\rho \right) \end{aligned} \quad (2.3)$$

Here, the interface couplings are all real functions of x^π . The couplings y_2^{ij} , y_3^{ijk} , z_2^{ij} , and z_3^{ijk} are totally anti-symmetric in their indices, while z_1^{ij} , and z_4^{ij} are totally symmetric. Overall factors of 2 and g have been introduced for later convenience. The following abbreviated notations will be useful,

$$\begin{aligned} Y_2 &\equiv y_2^{ij} \rho^{ij} & Y_3 &\equiv -y_3^{ijk} (\rho^{ijk})^* \\ Z_2 &\equiv z_2^{ij} \rho^{ij} & Z_3 &\equiv -z_3^{ijk} (\rho^{ijk})^* \end{aligned} \quad (2.4)$$

By construction, and using (A.5) of the Appendix, we have $Y_2^\dagger = -Y_2$, $Z_2^\dagger = -Z_2$, $Y_3^t = Y_3$, $Z_3^t = Z_3$, and $\text{tr}(Y_2) = \text{tr}(Z_2) = 0$. Under $SU(4)$, the interface operators and couplings transform in the following representations,

$$\begin{aligned} y_1 &: \mathbf{1} & z_1 &: \mathbf{1} \oplus \mathbf{20}' & z_4 &: \mathbf{1} \oplus \mathbf{20}' \\ y_2 &: \mathbf{15} & z_2 &: \mathbf{15} & z_{CS} &: \mathbf{1} \\ y_3 &: \mathbf{10} \oplus \overline{\mathbf{10}} & z_3 &: \mathbf{10} \oplus \overline{\mathbf{10}} \end{aligned} \quad (2.5)$$

In the context of AdS/CFT, the $\mathbf{10}$, $\overline{\mathbf{10}}$, $\mathbf{15}$, and $\mathbf{20}'$, correspond to supergravity fields, while the $\mathbf{1}$ corresponds to higher string modes. We do not expect the latter to be relevant in supergravity solutions, but shall include them here for completeness. Since we omitted the CP -violating term $\text{tr}(F\tilde{F})$ from $\mathbb{L}_0$, we shall omit its CP -violating interface counterpart $\mathbb{L}_{CS}$.

The final interface contributions to the Lagrangian of interest to us will be,

$$\mathbb{L}_I = \mathbb{L}_\psi + \mathbb{L}_\phi + \mathbb{L}_{\phi^2} \quad (2.6)$$

It is clear that all interface terms in $\mathbb{L}_\psi$, $\mathbb{L}_\phi$ correspond to local gauge invariant, dimension 3 operators whose contribution will be invariant under $SO(2, 3)$ conformal transformations in the limit where g tends to a step function.

It remains to establish that also the converse is true, namely that we have indeed retained all allowed operators. Combining gauge invariance and 2+1-dimensional Poincaré invariance with the requirement that the dimension of the interface operators can be at most 3, it is clear that no operators of dimension 1 can occur (since $\text{tr}(\phi^i) = 0$), and that the only dimension 2 operator is $\mathbb{L}_{\phi^2}$. Dimension 3 operators can then be (1) bilinears in the Fermi field, (2) trilinears in the scalar field or (3) bilinears in the scalar field with one extra derivative acting. $\mathbb{L}_\phi$ contains all possible terms in groups (2) and (3). The term $\text{tr}(\phi^i\{\phi^j, \phi^k\})$, which is totally symmetric in i, j, k , and which is a candidate for group (2), actually vanishes since the field ϕ^i takes values in the adjoint representation of the gauge algebra of $SU(N)$, (which is real) so that $\text{tr}(\phi^i\{\phi^j, \phi^k\}) = 0$. Finally, two types of fermion bilinears are allowed by $SO(1, 2)$ Lorentz invariance, and the fact that ψ is a Weyl spinor, namely $\text{tr}(\bar{\psi}\gamma^\pi M_1\psi)$ and $\text{tr}(\psi^t \mathcal{C} M_2 \psi)$ plus its complex conjugate. Here, M_1 and M_2 are 4×4 Clebsch-Gordan matrices of $SU(4)$, which in view of CPT -invariance and transposition symmetry of the bilinears must satisfy $M_1^\dagger = M_1$ and $M_2^t = M_2$. The unique solutions of these requirements are the ones listed in $\mathbb{L}_\psi$.

The interface couplings y_2^{ij} and z_2^{ij} may be viewed as external $SU(4) \sim SO(6)$ gauge fields which are localized at the interface, while y_1 may be viewed as an external $U(1)$ gauge field. Changing the fields ϕ^i and ψ by a local $SU(4)$ transformation which depends only on x^π will have the effect of making a gauge transformation on y_2^{ij} , z_2^{ij} , and z_4^{ij} . These transformations are free of anomalies and their existence shows that a given physical theory may be represented via different sets of interface couplings, as long as they are related by such an $SU(4)$ gauge transformation.

In the formulation of interface Lagrangians given above, it is straightforward to recover the two interface CFTs proposed in [11] as duals to the Janus solution. The first Lagrangian is obtained by choosing $z_1^{ij} = \delta^{ij}$ and all other parameters to be zero, while the second one is recovered by also choosing $z_4^{ij} = -\delta^{ij}$ and then redefining the scalar fields by $\phi^i \rightarrow \phi^i/g$. Interestingly, the $(\partial_\pi g)^2$ term now drops out for the redefined scalar fields, so that the limit to a step function gauge coupling is well defined. We shall list the general conditions for the existence of the conformal limit later.

3 Supersymmetry in the presence of an interface

In this section, we discuss the fate of supersymmetry in the interface Yang-Mills theory governed by the Lagrangian $\mathbb{L}$ of (2.2) whose interface operators and couplings are given in (2.3) and (2.6). We reduce the conditions for the existence of supersymmetry to a set of algebraic equations involving the gauge coupling, the interface couplings and the supersymmetry parameter. These equations will be solved in subsequent sections.

The supersymmetry transformations for the Lagrangian $\mathbb{L}_0$ are as follows [19, 20],²

$$\begin{aligned}\delta_0 A_\mu &= i\bar{\psi}\gamma_\mu\zeta - i\bar{\zeta}\gamma_\mu\psi \\ \delta_0 \phi^i &= i\zeta^t \mathcal{C} \rho^i \psi + i\bar{\psi} \mathcal{B}(\rho^i)^* \zeta^* \\ \delta_0 \psi &= +\frac{1}{2} F_{\mu\nu} \gamma^{\mu\nu} \zeta + (D_\mu \phi^i) \gamma^\mu \mathcal{B}(\rho^i)^* \zeta^* - \frac{i}{2} [\phi^i, \phi^j] \rho^{ij} \zeta\end{aligned}\quad (3.1)$$

It will be useful to have the transformations of the following fields as well,

$$\begin{aligned}\delta_0 F_{\mu\nu} &= iD_\mu \bar{\psi} \gamma_\nu \zeta - i\bar{\zeta} \gamma_\nu D_\mu \psi - iD_\nu \bar{\psi} \gamma_\mu \zeta + i\bar{\zeta} \gamma_\mu D_\nu \psi \\ \delta_0 (D_\mu \phi^i) &= i\zeta^t \mathcal{C} \rho^i D_\mu \psi + i\zeta^\dagger \mathcal{C}(\rho^i)^* D_\mu \psi^* + [\bar{\psi} \gamma_\mu \zeta - \bar{\zeta} \gamma_\mu \psi, \phi^i] \\ \delta_0 \bar{\psi} &= -\frac{1}{2} \bar{\zeta} \gamma^{\mu\nu} F_{\mu\nu} - (D_\mu \phi^i) \zeta^t \mathcal{C} \rho^i \gamma^\mu + \frac{i}{2} [\phi^i, \phi^j] \bar{\zeta} \rho^{ij}\end{aligned}\quad (3.2)$$

Under the transformation δ_0 , with space-time dependent supersymmetry parameter ζ , the Lagrangian transforms as follows,³

$$\delta_0 \mathbb{L}_0 = \frac{1}{g^2} \left(\partial_\kappa X^\kappa - (\partial_\kappa \bar{\zeta}) S^\kappa \right) \quad (3.3)$$

where

$$\begin{aligned}X^\kappa &= \frac{i}{4} \text{tr} \left(F_{\mu\nu} \bar{\psi} \gamma^{\mu\nu} \gamma^\kappa \zeta - 2D_\mu \phi^i \bar{\psi} \gamma^\mu \gamma^\kappa \mathcal{B}(\rho^i)^* \zeta^* - i[\phi^i, \phi^j] \bar{\psi} \gamma^\kappa \rho^{ij} \zeta \right) + \text{c.c.} \\ (\partial_\kappa \bar{\zeta}) S^\kappa &= \frac{i}{2} \text{tr} \left(F_{\mu\nu} \bar{\psi} \gamma^\kappa \gamma^{\mu\nu} \partial_\kappa \zeta + 2D_\mu \phi^i \bar{\psi} \gamma^\kappa \gamma^\mu \mathcal{B}(\rho^i)^* \partial_\kappa \zeta^* - i[\phi^i, \phi^j] \bar{\psi} \gamma^\kappa \rho^{ij} \partial_\kappa \zeta \right) + \text{c.c.}\end{aligned}\quad (3.4)$$

For constant g , it is clear from the transformation (3.3), that the Lagrangian $\mathbb{L}_0$ is invariant under the global symmetry generated by δ_0 with constant ζ , for which S^κ is the conserved supercurrent. Indeed, for constant g and ζ , the Lagrangian changes by a total derivative $\partial_\kappa (g^{-2} X^\kappa)$, signaling that δ_0 is a symmetry.

²The complex conjugation matrix B is defined to obey the relation $\mathcal{B} \gamma_\mu \mathcal{B}^{-1} = (\gamma_\mu)^*$; it may be chosen so that $\mathcal{B}^* = \mathcal{B}$, $\mathcal{B}^2 = I$, and is given by $\mathcal{B} = \mathcal{C} \gamma^0$.

³The customary Majorana notation $(\partial_\kappa \bar{\zeta}) S^\kappa$ includes the contribution from ζ and ζ^* .

3.1 Conditions for interface supersymmetry

For non-constant g , (such as when g is a function of x^π) the Lagrangian $\mathbb{L}_0$ no longer changes by a total derivative under the transformation δ_0 with constant ζ , but instead we have now the following transformation equation,

$$\delta_0 \mathbb{L}_0 = \partial_\kappa \left(\frac{1}{g^2} X^\kappa \right) - \frac{1}{g^2} (\partial_\kappa \bar{\zeta}) S^\kappa + 2 \frac{\partial_\kappa g}{g^3} X^\kappa \quad (3.5)$$

As a result, supersymmetry will be broken. For an interface described by a gauge coupling $g(x^\pi)$, the obstructing term is proportional to $\partial_\pi g$ and is thus localized on the interface. Therefore, it is natural to modify the Lagrangian by including the interface operators of (2.3) and (2.6), to modify the supersymmetry transformations by interface terms δ_I proportional to $\partial_\pi g$,

$$\delta = \delta_0 + \delta_I \quad (3.6)$$

and to let the supersymmetry parameter depend on x^π . Restricting the form of δ_I by 2+1-dimensional Poincaré invariance, gauge invariance and the requirement that no extra dimensionful parameters should enter, we find that,

$$\begin{aligned} \delta_I \phi^i &= \delta_I A_\mu = 0 \\ \delta_I \psi &= (\partial_\pi g) \chi^i \phi^i \end{aligned} \quad (3.7)$$

The spinors⁴ χ^i have scaling dimension $-1/2$, and are as of yet undetermined functions of x^π . The scaling dimension is due to a factor of ζ , which will be shown later (3.15), so that χ^i does not introduce any dimensionful parameters. The requirement of supersymmetry of the full Lagrangian $\mathbb{L}$ of (2.2) under the full transformations δ of (3.6) is that the full variation,

$$\delta \mathbb{L} = \delta_0 \mathbb{L}_0 + \delta_0 \mathbb{L}_I + \delta_I \mathbb{L}_0 + \delta_I \mathbb{L}_I \quad (3.8)$$

be a total derivative. Since g and ζ are smooth functions of only the single space-coordinate x^π , this condition reduces to

$$-\frac{1}{g^2} (\partial_\pi \bar{\zeta}) S^\pi + 2 \frac{\partial_\pi g}{g^3} X^\pi + \delta_0 \mathbb{L}_I + \delta_I \mathbb{L}_0 + \delta_I \mathbb{L}_I \sim 0 \quad (3.9)$$

up to a total derivative. The calculations of $\delta_0 \mathbb{L}_I$, $\delta_I \mathbb{L}_0$, and $\delta_I \mathbb{L}_I$ are straightforward, but the resulting expressions are lengthy and will not be presented here.

⁴ χ^i is a Weyl spinor, in the **4** of $SU(4)$, and is a singlet under the $SU(N)$ gauge symmetry.

Besides total derivative terms, the equation (3.9) involves different functionally independent combinations of the canonical fields ϕ^i , ψ , $\bar{\psi}$, and A_μ , which must vanish independently. Half of these terms involve ψ , the other half involve $\bar{\psi}$, which are complex conjugates of one another. It suffices to enforce the vanishing of the terms in $\bar{\psi}$, as the ensuing equations will, by complex conjugation, also imply the vanishing of the terms in ψ . The independent field combinations are $\phi^i\bar{\psi}$, $\phi^i D_\pi \bar{\psi}$, $F_{\mu\nu}\bar{\psi}$, $(D_\mu \phi^i)\bar{\psi}$, and $[\phi^j, \phi^k]\bar{\psi}$.⁵ Their coefficients must independently vanish, and yield the following equations;⁶

- The coefficients of $\phi^i\bar{\psi}$ are,

$$0 = (-iy_1 + \frac{1}{4}Y_2 - 1)\gamma^\pi \chi^i - \mathcal{B}Y_3(\chi^i)^* + z_4^{ij}\mathcal{B}(\rho^j)^*\zeta^* + (z_1^{ij} + z_2^{ij})\mathcal{B}(\rho^j)^*\frac{\partial_\pi \zeta^*}{\partial_\pi g} \quad (3.10)$$

- The coefficients of $\phi^i D_\pi \bar{\psi}$ are,

$$0 = (z_1^{ij} + z_2^{ij})\mathcal{B}(\rho^j)^*\zeta^* + \gamma^\pi \chi^i \quad (3.11)$$

- The coefficients of $F_{\mu\nu}\bar{\psi}$ are,

$$0 = \gamma^{\mu\nu}\gamma^\pi \zeta - \gamma^\pi \gamma^{\mu\nu} \frac{\partial_\pi \zeta}{\partial_\pi g} - iy_1 \gamma^\pi \gamma^{\mu\nu} \zeta + \frac{1}{4} \gamma^\pi \gamma^{\mu\nu} Y_2 \zeta - \gamma^{\mu\nu} \mathcal{B}Y_3 \zeta^* \quad (3.12)$$

- The coefficients of $\bar{\psi}(2D_\mu \phi^i)$ are,

$$\begin{aligned} 0 = & -\gamma^\mu \gamma^\pi \mathcal{B}(\rho^i)^*\zeta^* - \gamma^\pi \gamma^\mu \mathcal{B}(\rho^i)^*\frac{\partial_\pi \zeta^*}{\partial_\pi g} - iy_1 \gamma^\pi \gamma^\mu \mathcal{B}(\rho^i)^*\zeta^* + \frac{1}{4} \gamma^\pi \gamma^\mu \mathcal{B}Y_2(\rho^i)^*\zeta^* \\ & + \gamma^\mu Y_3 \rho^i \zeta - z_2^{ij} \mathcal{B}(\rho^j)^*\zeta^* \delta^{\mu\pi} + z_1^{ij} \mathcal{B}(\rho^j)^*\zeta^* \delta^{\mu\pi} - \gamma^\mu \chi^i + \gamma^\pi \chi^i \delta^{\mu\pi} \end{aligned} \quad (3.13)$$

- The coefficients of $i[\phi^i, \phi^j]\bar{\psi}$ are,

$$\begin{aligned} 0 = & -\gamma^\pi \rho^{ij} \zeta + \gamma^\pi \rho^{ij} \frac{\partial_\pi \zeta}{\partial_\pi g} + iy_1 \gamma^\pi \rho^{ij} \zeta - \frac{1}{4} \gamma^\pi Y_2 \rho^{ij} \zeta + \mathcal{B}Y_3(\rho^{ij})^*\zeta^* \\ & - 3z_3^{ijk} \mathcal{B}(\rho^k)^*\zeta^* + 2z_2^{ij} \gamma_\pi \zeta + \mathcal{B}(\rho^i)^*(\chi^j)^* - \mathcal{B}(\rho^j)^*(\chi^i)^* \end{aligned} \quad (3.14)$$

In (3.10), we have omitted a common factor of $i(\partial_\pi g)^2$, while in (3.11-3.14), we have omitted a common factor of $i(\partial_\pi g)/2$. Finally, all terms with no $\partial_\pi g$ dependence are total derivatives and do not contribute to the variation of the action.

⁵Note that $D_\pi(\phi\bar{\psi})$ will not yield a total derivative in the action so that $\phi^i D_\pi \bar{\psi}$ and $(D_\mu \phi^i)\bar{\psi}$ are functionally independent.

⁶Throughout, we continue to use the notations of (2.4) for Y_2 and Y_3 .

3.2 Simplifying the Conditions for Interface Supersymmetry

We shall now simplify and to some extent decouple the equations for interface supersymmetry, obtained in (3.10-3.14). The result is the following group of 7 equations,

$$\begin{aligned}
(1) \quad & \zeta = Y_3 \gamma^\pi \mathcal{B} \zeta^* \\
(2) \quad & Y_3 \rho^i \zeta = -(\rho^i)^* \gamma^\pi \mathcal{B} \zeta^* + z_1^{ij} (\rho^j)^* \gamma^\pi \mathcal{B} \zeta^* \\
(3) \quad & -\rho^{ij} \zeta + y_1 i \rho^{ij} \zeta + \rho^{ij} \frac{\partial_\pi \zeta}{\partial_\pi g} - \frac{1}{4} Y_2 \rho^{ij} \zeta + Y_3 (\rho^{ij})^* \gamma^\pi \mathcal{B} \zeta^* - 3 z_3^{ijk} (\rho^k)^* \gamma^\pi \mathcal{B} \zeta^* \\
& + 2 z_2^{ij} \zeta - (\rho^i)^* (z_1^{jk} + z_2^{jk}) \rho^k \zeta + (\rho^j)^* (z_1^{ik} + z_2^{ik}) \rho^k \zeta = 0 \\
(4) \quad & \frac{\partial_\pi \zeta}{\partial_\pi g} = -i y_1 \zeta + \frac{1}{4} Y_2 \zeta \\
(5) \quad & \rho^i \frac{\partial_\pi \zeta}{\partial_\pi g} = i y_1 \rho^i \zeta + \frac{1}{4} Y_2^* \rho^i \zeta - z_2^{ij} \rho^j \zeta \\
(6) \quad & \chi^i = -(z_1^{ij} + z_2^{ij}) (\rho^j)^* \gamma^\pi \mathcal{B} \zeta^* \\
(7) \quad & (-i y_1 + \frac{1}{4} Y_2 - 1) (z_1^{ij} + z_2^{ij}) (\rho^j)^* \gamma^\pi \mathcal{B} \zeta^* - Y_3 (z_1^{ij} + z_2^{ij}) \rho^j \zeta \\
& - z_4^{ij} (\rho^j)^* \gamma^\pi \mathcal{B} \zeta^* - (z_1^{ij} + z_2^{ij}) (\rho^j)^* \frac{\partial_\pi \gamma^\pi \mathcal{B} \zeta^*}{\partial_\pi g} = 0
\end{aligned} \tag{3.15}$$

These equations were obtained as follows. Multiplying (3.11) by γ^π yields equation (6), which now gives χ^i in terms of the other variables. Equation (6) may be used to eliminate χ^i from all other equations; doing so in (3.10) yields equation (7), while doing so in (3.14) yields equation (3).

To simplify (3.12), we use the fact that, as μ and ν run through their possible values 0, 1, 2, 3, the Clifford generators $\gamma^{\mu\nu} \gamma^\pi$ are generally linearly independent from $\gamma^\pi \gamma^{\mu\nu}$, so that their coefficients in (3.12) must vanish independently; this yields equations (1) and (4) respectively. Equivalently, one may consider the two distinct cases $\pi \in \{\mu, \nu\}$ and $\pi \notin \{\mu, \nu\}$; equations (1) and (4) are then obtained as the sum and difference of these two conditions.

Similarly, to simplify (3.13), we use the fact that, as μ run through 0, 1, 2, 3, the Clifford generators $\gamma^\mu \gamma^\pi$ are generally linearly independent from $\gamma^\pi \gamma^\mu$, so that their coefficients in (3.13) must vanish independently. Using in these equations also (6) to eliminate χ^i , yields (2) and (5) respectively, after some simplifications. Equivalently, one may consider the two distinct cases $\pi = \mu$ and $\pi \neq \mu$; equations (2) and (5) are then obtained as the sum and difference of these two conditions.

3.3 Resolving the space-time spinor structure

The preceding equations involve the spinors ζ and $\gamma^\pi \mathcal{B} \zeta^*$ with coefficients which act on the internal $SU(4)$ labels of the spinors, but not on their 3+1-dimensional space-time spinor labels. Clearly, it will be convenient to decompose the spinors ζ and $\gamma^\pi \mathcal{B} \zeta^*$ onto a basis in which $\gamma^\pi \mathcal{B}$ is diagonal. This is achieved in this subsection, and the reduced equations are then derived. For definiteness, we choose $\mathcal{B}$ to obey the relations of footnote #2, namely $\mathcal{B}^* = \mathcal{B}^t = \mathcal{B}^{-1} = \mathcal{B}$, so that we have

$$(\gamma^\pi \mathcal{B})^t = \gamma^\pi \mathcal{B} \quad (\gamma^\pi \mathcal{B})^* = \mathcal{B} \gamma^\pi = (\gamma^\pi \mathcal{B})^{-1} \quad (3.16)$$

As a result, the matrix $\gamma^\pi \mathcal{B}$ is unitary, and has eigenvalues of unit modulus. Furthermore, $\gamma^\pi \mathcal{B}$ commutes with the 3+1-dimensional chirality matrix γ . Finally, $\det(\gamma^\pi \mathcal{B}) = 1$, while on the $+$ chirality subspace, we have $\det(\gamma^\pi \mathcal{B})_+ = -1$.

Using the above facts about $\gamma^\pi \mathcal{B}$, we conclude that, on the $+$ chirality subspace, $\gamma^\pi \mathcal{B}$ has two eigenvalues, λ^2 and $-(\lambda^2)^*$, with $\lambda \lambda^* = 1$. Since $\gamma^\pi \mathcal{B}$ is unitary, it is diagonalizable, and we denote its eigenvectors by $s_\pm$,

$$\begin{aligned} \gamma^\pi \mathcal{B} s_+ &= +\lambda^2 s_+ \\ \gamma^\pi \mathcal{B} s_- &= -(\lambda^2)^* s_- \end{aligned} \quad (3.17)$$

Using $(\gamma^\pi \mathcal{B})^* = \mathcal{B} \gamma^\pi$ and $\lambda \lambda^* = 1$, it follows that

$$\begin{aligned} \gamma^\pi \mathcal{B} s_+^* &= +\lambda^2 s_+^* \\ \gamma^\pi \mathcal{B} s_-^* &= -(\lambda^2)^* s_-^* \end{aligned} \quad (3.18)$$

When the two eigenvalues are different ($\text{Re}(\lambda^2) \neq 0$), the eigenspaces are 1-dimensional, and we may choose $s_+^* = s_+$ and $s_-^* = s_-$. When $\text{Re}(\lambda^2) = 0$, $\gamma^\pi \mathcal{B}$ is proportional to the identity and the same choice is still valid. We now decompose ζ and $\partial_\pi \zeta$ as follows,

$$\begin{aligned} \zeta &= \lambda s_+ \otimes \xi_+ + i\lambda^* s_- \otimes \xi_- \\ \partial_\pi \zeta &= \lambda s_+ \otimes \partial_\pi \xi_+ + i\lambda^* s_- \otimes \partial_\pi \xi_- \end{aligned} \quad (3.19)$$

Applying the above rules for complex conjugation, we then derive,

$$\begin{aligned} \gamma^\pi \mathcal{B} \zeta^* &= \lambda s_+ \otimes \xi_+^* + i\lambda^* s_- \otimes \xi_-^* \\ \gamma^\pi \mathcal{B} \partial_\pi \zeta^* &= \lambda s_+ \otimes \partial_\pi \xi_+^* + i\lambda^* s_- \otimes \partial_\pi \xi_-^* \end{aligned} \quad (3.20)$$

3.4 The reduced conditions for interface supersymmetry

The equations of (3.15) may be decomposed onto the basis of (3.19) and (3.20). It is manifest that both ξ_+ and $\partial_\pi \xi_+$, and ξ_- and $\partial_\pi \xi_-$ satisfy the same equations. Dropping the $\pm$ subscripts on ξ and $\partial_\pi \xi$, we get

$$\begin{aligned}
(1') \quad & \xi = Y_3 \xi^* \\
(2') \quad & Y_3 \rho^i \xi = -(\rho^i)^* \xi^* + z_1^{ij} (\rho^j)^* \xi^* \\
(3') \quad & \left(\rho^{ij} Y_3 + Y_3 (\rho^{ij})^* - (\rho^i)^* Y_3^* (\rho^j)^* + (\rho^j)^* Y_3^* (\rho^i)^* \right) \xi^* - 3 z_3^{ijk} (\rho^k)^* \xi^* \\
& + \frac{1}{4} \left(\rho^{ij} Y_2 - Y_2 \rho^{ij} \right) \xi + 2 z_2^{ij} \xi - (\rho^i)^* z_2^{jk} \rho^k \xi + (\rho^j)^* z_2^{ik} \rho^k \xi = 0 \\
(4') \quad & \partial_\pi \xi = (\partial_\pi g) \left(-i y_1 \xi + \frac{1}{4} Y_2 \xi \right) \\
(5') \quad & 2 i y_1 \rho^i \xi + \frac{1}{4} \left(Y_2^* \rho^i - \rho^i Y_2 \right) \xi - z_2^{ij} \rho^j \xi = 0 \\
(6) \quad & \chi^i = - \left(z_1^{ij} + z_2^{ij} \right) (\rho^j)^* \gamma^\pi \mathcal{B} \zeta^* \\
(7') \quad & \left(z_4^{ij} + (z_1^{ik} + z_2^{ik})(z_1^{jk} + z_2^{jk}) \right) (\rho^j)^* \xi^* = 0
\end{aligned} \tag{3.21}$$

We have used (4') to eliminate $\partial_\pi \xi$ from the equations leading to (3'), (5'), and (7'); used (2') to eliminate z_1^{ij} from the equation leading to (3'), and $Y_3 \rho^j \xi$ from the equation leading to (7'); used (5') to further simplify the equation that leads to (7'). Finally, (6) has been left undecomposed, as it merely gives an expression for χ^i . For every ξ satisfying these equations, we will have two real supersymmetries given by ξ_+ and ξ_- .

Given the assumption that a non-trivial supersymmetry survive requires solutions with $\xi \neq 0$. This in turn allows for a simple solution to equation (7')

$$(7') \quad z_4^{ij} = -(z_1^{ik} + z_2^{ik})(z_1^{jk} + z_2^{jk}) \tag{3.22}$$

Since the variable z_4^{ij} does not appear in the remaining equations, we can take (3.22) as the resulting value of z_4^{ij} .

4 Solving the Interface Supersymmetry Conditions

In this section, we shall solve the conditions for interface supersymmetry which, in the preceding section, were reduced to a set of algebraic equations (3.21). To simplify the associated calculations, we shall use the covariance of the interface couplings, supersymmetry parameter, and equations under global and local $SU(4)$ transformations.

4.1 Transformations under global $SU(4)$

Let U and R be the representation matrices respectively in the **4** and **6** of $SU(4)$, satisfying $U^\dagger U = I$, $R^t R = I$. The fields and supersymmetry parameters transform as

$$\begin{aligned}\psi &\rightarrow \psi' = U\psi & \zeta &\rightarrow \zeta' = U\zeta \\ \phi^i &\rightarrow (\phi')^i = R^{ii'} \phi^{i'} & \xi &\rightarrow \xi' = U\xi\end{aligned}\tag{4.1}$$

The ρ -matrices are invariant in the following sense,⁷

$$\begin{aligned}\rho^i &= R^{ii'} U^* \rho^{i'} U^\dagger \\ \rho^{ij} &= R^{ii'} R^{jj'} U \rho^{i'j'} U^\dagger \\ \rho^{ijk} &= R^{ii'} R^{jj'} R^{kk'} U^* \rho^{i'j'k'} U^\dagger\end{aligned}\tag{4.2}$$

Under the above $SU(4)$ transformations, L_0 is invariant, even with varying gauge coupling $g(x^\pi)$. In general, however, the interface Lagrangian L_I is not invariant because the interface couplings transform non-trivially. Thus, $SU(4)$ maps one interface theory onto another by mapping their respective interface couplings,

$$\begin{aligned}y^{i_1 \dots i_n} &\rightarrow (y')^{i_1 \dots i_n} = R^{i_1 i'_1} \dots R^{i_n i'_n} y^{i'_1 \dots i'_n} \\ z^{i_1 \dots i_n} &\rightarrow (z')^{i_1 \dots i_n} = R^{i_1 i'_1} \dots R^{i_n i'_n} z^{i'_1 \dots i'_n}\end{aligned}\tag{4.3}$$

but leaving the gauge coupling unchanged. Under these combined transformations, the reduced equations (3.21) are also invariant. Combining (4.2) with (4.3), we have the following convenient formulas,

$$\begin{aligned}Y_2 &\rightarrow Y'_2 = U Y_2 U^\dagger & Y_3 &\rightarrow Y'_3 = U Y_3 U^t \\ Z_2 &\rightarrow Z'_2 = U Z_2 U^\dagger & Z_3 &\rightarrow Z'_3 = U Z_3 U^t\end{aligned}\tag{4.4}$$

⁷This result may be derived by starting from the $SO(6)$ Dirac matrices $\gamma_{(6)}^i$, for which we have $\gamma_{(6)}^i = R^{ii'} U \gamma_{(6)}^{i'} U^\dagger$ and $U^* = \mathcal{C}_{(6)} U \mathcal{C}_{(6)}^{-1}$, and using the relation (A.3).

4.2 Further simplifying (3') and (5')

To exhibit covariance of (3.21), (3.22) and (4.5) under local x^π -dependent $SU(4)$ transformations, and then solve these equations, we shall begin by further simplifying the form of (3') and (5'). To do so, we express Y_3 in terms of its coefficients y_3^{ijk} , and z_2^{ij} in terms of the matrix Z_2 , both using (2.4),

$$\begin{aligned} (3') \quad & 3 \left(z_3^{ijk} - 8y_3^{ijk} \right) (\rho^k)^* \xi^* - \frac{1}{4} \left(\rho^{ij} (Y_2 + Z_2) - (Y_2 + Z_2) \rho^{ij} \right) \xi = 0 \\ (5') \quad & 2iy_1 \rho^i \xi + \frac{1}{4} \left((Y_2 + Z_2)^* \rho^i - \rho^i (Y_2 + Z_2) \right) \xi = 0 \end{aligned} \quad (4.5)$$

using the relations,

$$\begin{aligned} Z_2^* \rho^i - \rho^i Z_2 &= -4z_2^{ij} \rho^j \\ \rho^{ij} Z_2 - Z_2 \rho^{ij} &= 8z_2^{ij} - 4z_2^{jk} (\rho^i)^* \rho^k + 4z_2^{ik} (\rho^j)^* \rho^k \\ 24y_3^{ijk} (\rho^k)^* &= \rho^{ij} Y_3 + Y_3 (\rho^{ij})^* - (\rho^i)^* Y_3^* (\rho^j)^* + (\rho^j)^* Y_3^* (\rho^i)^* \end{aligned} \quad (4.6)$$

The first two equations above were derived by contracting identities (A.8) and (A.9) with z_2^{kl} , and using (A.7) to rearrange the second equation in the form above. The last equation was derived by contracting (A.10) with y_3^{klm} , and making further simplifications using (A.7) and (A.8).

4.3 Transformations under local $SU(4)$

The fact that the interface couplings y_2^{ij} and z_2^{ij} may be regarded as arising from a $SU(4)$ background gauge field implies that the Lagrangian $\mathcal{L}$, and the interface supersymmetry conditions (3.21) behave covariantly under local x^π -dependent $SU(4)$ transformations.

Considering the equations of (3.21), and their simplified forms in (3.22) and (4.5), we see that only (4') involves a derivative with respect to x^π . Thus, equations (1'), (2'), (3'), (5') and (7') are actually covariant under *local* $SU(4)$ transformations $\mathcal{U}$, and associated $\mathcal{R}$, whose action is defined as follows,

$$\begin{aligned} \xi &\rightarrow \xi' = \mathcal{U} \xi \\ \rho^i &= \mathcal{R}^{ii'} \mathcal{U}^* \rho^{i'} \mathcal{U}^\dagger \end{aligned} \quad (4.7)$$

Covariance of (1') and (2') then requires the following transformation laws,

$$\begin{aligned} Y_3 &\rightarrow Y'_3 = \mathcal{U} Y_3 \mathcal{U}^t \\ z_1^{ij} &\rightarrow (z'_1)^{ij} = \mathcal{R}^{ii'} \mathcal{R}^{jj'} z_1^{i'j'} \end{aligned} \quad (4.8)$$

Covariance of the differential equation (4') requires that Y_2 transform as a $SU(4)$ connection, while (5') requires that $Y_2 + Z_2$ transform homogeneously. As a result,

$$\begin{aligned} Y_2 &\rightarrow Y'_2 = \mathcal{U} Y_2 \mathcal{U}^\dagger + \omega & \omega &= \frac{4}{\partial_\pi g} (\partial_\pi \mathcal{U}) \mathcal{U}^\dagger \\ Z_2 &\rightarrow Z'_2 = \mathcal{U} Z_2 \mathcal{U}^\dagger - \omega \end{aligned} \quad (4.9)$$

and this induces the following transformation on z_4^{ij} ,

$$z_4^{ij} \rightarrow (z'_4)^{ij} = \mathcal{R}^{ii'} \mathcal{R}^{jj'} \left(z_4^{ij} + \omega^{ik} (z_1^{jk} + z_2^{jk}) + \omega^{jk} (z_1^{ik} + z_2^{ik}) - \omega^{ik} \omega^{jk} \right) \quad (4.10)$$

Under the above transformations, the equations (3.21), and their simplified forms in (3.22) and (4.5), are covariant.

The $SU(4)$ gauge transformation, introduced above, is free of anomalies and thus remains a viable map between full quantum theories. This may be seen as follows. Its mixed anomalies with any semi-simple gauge current vanish as both the $SU(4)$ and the gauge generators are traceless. The triangle graph with three $SU(4)$ currents vanishes for kinematic reasons. This may be seen from the fact that the only non-vanishing components of the “gauge field” is $A_\pi(x^\pi) \sim y_2$ for which $\epsilon^{\mu\nu\rho\sigma} A_\nu \partial_\rho A_\sigma$ and $\epsilon^{\mu\nu\rho\sigma} A_\nu A_\rho A_\sigma$ vanish identically, thereby canceling the perturbative anomaly.

4.4 Choosing the gauge $Y_2 = 0$ and solving (4')

The covariance under local $SU(4)$, established in the preceding section, allows us to choose $\mathcal{U}(x^\pi)$ at will. Different choices of $\mathcal{U}$ will lead to different values for the interface couplings, but to physically equivalent field theories. It will be convenient to choose

$$Y_2 = 0 \quad (4.11)$$

since the differential equation may then be easily solved,

$$\begin{aligned} \xi(x^\pi) &= \xi_0 \beta e^{i\theta(x^\pi)/2} \\ \theta(x^\pi) &= \theta(x_0^\pi) - 2 \int_{x_0^\pi}^{x^\pi} dx^\pi (\partial_\pi g) y_1(x^\pi) \end{aligned} \quad (4.12)$$

Here, ξ_0 is a real, constant, Grassmann number; β is a real constant 4-vector, left undetermined by equation (4'); and x_0^π is an arbitrary reference point. It will be important in the sequel that ξ maintains a direction independent of x^π .

4.5 Diagonalizing Y_3

From equation (1') in (3.21), it is manifest that $\text{rank}(I - Y_3^* Y_3)$ provides an upper bound on the number of allowed interface supersymmetries. In particular, when $Y_3 = 0$, as is the case for the CFT dual of the Janus solution, no interface supersymmetries will exist. Of course, even when $\text{rank}(I - Y_3^* Y_3) > 1$, the solutions to (1') will also have to satisfy (2'), (3') and (5'), which will impose further restrictions. In this subsection, we use global $SU(4)$ transformations to solve (1') completely, as a function of Y_3 .

We proceed as follows. We work first at one arbitrary point x_0^π . Using only global $SU(4)$, we simplify (3.21) by choosing a convenient frame to express the interface couplings. This simplification is carried out by first diagonalizing Y_3 with the help of the following little

Lemma 1 A complex symmetric matrix $Y_3 = Y_3^t$ admits the following diagonalization,

$$Y_3 = U D_3 U^t e^{i\theta_0} \quad (4.13)$$

where D_3 is real diagonal, with non-negative entries, θ_0 is a real phase, and U is special unitary, satisfying $U^\dagger U = I$, and $\det U = 1$. (When $\det(D_3) = 0$, the angle θ_0 may be absorbed in the special unitary rotation U .) Note that the number of independent parameters on both sides are equal: 20 for Y_3 ; 15 for U ; 4 for D_3 , and 1 for θ_0 .

We review the proof here. Using $Y_3^t = Y_3$, the matrix $Y_3 Y_3^* = Y_3 Y_3^\dagger$ is seen to be Hermitian and non-negative, and may be diagonalized by a special unitary matrix U ,

$$Y_3 Y_3^* = U D_3^2 U^\dagger \quad (4.14)$$

Here, D_3 is real diagonal; its entries may be chosen to be non-negative. We prove the Lemma first when D_3 is invertible and has all distinct eigenvalues, in which case we have, $(D_3^{-1} U^\dagger Y_3) (D_3^{-1} U^\dagger Y_3)^\dagger = I$. The matrix $D_3^{-1} U^\dagger Y_3$ must be unitary (but not necessarily special unitary), and will be denoted by U_1 , so that $Y_3 = U D_3 U_1$. More generally,

$$Y_3 = U \Lambda D_3 \Lambda^* U_1 \quad (4.15)$$

where Λ is an arbitrary diagonal special unitary matrix. Equating the transpose of (4.14), $Y_3^* Y_3 = U^* D_3^2 U^t$, with $Y_3^\dagger Y_3 = U_1^\dagger \Lambda D_3^2 \Lambda^* U_1$, obtained from (4.15), we find that the matrix $\Lambda^* U_1 U^*$ must commute with D_3^2 . Since the eigenvalues of D_3^2 were assumed to be all distinct, $\Lambda^* U_1 U^*$ must be diagonal and unitary (though not necessarily special unitary). Since Λ is an arbitrary special unitary matrix, upon which U and U_1 do not depend, Λ may be chosen so that $U_1 U^* = I e^{i\theta_0}$, which proves Lemma 1 when D_3 is invertible and has all distinct eigenvalues.

When D_3 has zero and/or coincident eigenvalues, we make a small perturbation on Y_3 continuously parametrized by ε which preserves $Y_3^t = Y_3$ but lifts any degeneracies and moves all eigenvalues of D_3 away from 0. The Lemma now holds for small ε , of which D_3 , U and θ_0 are continuous functions. The limit $\varepsilon \rightarrow 0$ clearly exists since U and $e^{i\theta_0}$ take values in compact spaces. Finally, when $\det(D_3) = 0$, the matrix D_3 has at least one zero on the diagonal, the presence of which may be used to rotate away θ_0 (a mechanism well-known to those familiar with the strong CP-problem).

4.6 Solving equation (1')

We start from the parametrization $Y_3 = UD_3U^t e^{i\theta_0}$ proven in Lemma 1. Equation (1') has non-vanishing solutions ξ provided at least one eigenvalue of D_3 equals 1. Henceforth, we shall make this assumption, as we are only interested in interface theories with non-trivial supersymmetry. Without loss of generality, we may choose the ordering of the eigenvalues as follows,

$$D_3 = \text{diag}[1, a, b, c] \quad a, b, c \in \mathbf{R}^+ \quad (4.16)$$

We introduce a basis of eigenvectors β_a , $a = 1, 2, 3, 4$ with components $(\beta_a)_b = \delta_{ab}$. The existence of a non-vanishing solution ξ , whose x^π dependence was already determined in (4.12) from solving (4'), fixes the x^π -dependence of Y_3 .,

$$\begin{aligned} Y_3(x^\pi) &= UD_3U^t e^{i\theta(x^\pi)} \\ \xi(x^\pi) &= \xi_0 \beta e^{i\theta(x^\pi)/2} \quad \beta = U\beta_1 \end{aligned} \quad (4.17)$$

Since β was x^π -independent by (4.12), the $SU(4)$ matrix U must be x^π -independent as well. Therefore, U may be rotated away using only a global $SU(4)$ rotation. Whenever a , b , or c equals 1, more than a single solution will exist, with the same x^π -dependence but with β_1 replaced by a β_a , $a = 2, 3, 4$.

4.7 Solving equation (2')

Equation (2') now simplifies to give

$$e^{2i\theta} D_3 \rho^i \beta_1 = -(\rho^i)^* \beta_1 + z_1^{ij} (\rho^j)^* \beta_1 \quad (4.18)$$

The following simple relations, deduced from the explicit form of the matrices ρ^i in (A.13) of Appendix A,

$$\begin{aligned} \rho^1 \beta_1 &= i \beta_2 & \rho^3 \beta_1 &= -i \beta_3 & \rho^5 \beta_1 &= +i \beta_4 \\ \rho^2 \beta_1 &= +\beta_2 & \rho^4 \beta_1 &= -\beta_3 & \rho^6 \beta_1 &= +\beta_4 \end{aligned} \quad (4.19)$$

allow us to solve (4.18) by inspection, and we find,

$$\begin{aligned}
z_1^{11} &= 1 - a \cos(2\theta) & z_1^{12} &= z_1^{21} = -a \sin(2\theta) & z_1^{22} &= 1 + a \cos(2\theta) \\
z_1^{33} &= 1 - b \cos(2\theta) & z_1^{34} &= z_1^{43} = -b \sin(2\theta) & z_1^{44} &= 1 + b \cos(2\theta) \\
z_1^{55} &= 1 - c \cos(2\theta) & z_1^{56} &= z_1^{65} = -c \sin(2\theta) & z_1^{66} &= 1 + c \cos(2\theta)
\end{aligned} \quad (4.20)$$

with all other $z_1^{ij} = 0$.

4.8 Solving equation (5')

To solve (5') when $Y_2 = 0$, it will be convenient to use the original form of (5') in (3.21), and then substitute the solution (4.12) for ξ ,

$$2iy_1\rho^i\beta_1 - z_2^{ij}\rho^j\beta_1 = 0 \quad (4.21)$$

Using (4.19), this equation may also be solved by inspection. Clearly, all components of z_2^{ij} must vanish, except

$$z_2^{12} = z_2^{34} = z_2^{56} = -z_2^{21} = -z_2^{43} = -z_2^{65} = -2y_1 \quad (4.22)$$

It will be useful to recast this solution in the form of the matrix Z_2 , and we find,

$$Z_2 = -4iy_1 \left(I - 4\beta_1 \otimes \beta_1^\dagger \right) \quad (4.23)$$

4.9 Solving equation (3')

With the help of the preceding results for Y_2 , Y_3 , and Z_2 , equation (3') becomes an equation for z_3^{ijk} . Indeed, substituting $Y_2 = 0$ and the solution of (4.23) for Z_2 into (3') in (4.5), yields the following expression,

$$\left(z_3^{ijk} - 8y_3^{ijk} \right) (\rho^k)^* \beta_1 = \frac{4}{3} i y_1 e^{i\theta} \left(\rho^{ij} \beta_1 - \beta_1 \left(\beta_1^\dagger \rho^{ij} \beta_1 \right) \right) \quad (4.24)$$

Both sides are manifestly orthogonal to β_1 . Using again (4.19), the projections onto $\beta_2, \beta_3, \beta_4$ may be computed as follows,

$$\begin{aligned}
z_3^{ij1} - 8y_3^{ij1} + i \left(z_3^{ij2} - 8y_3^{ij2} \right) &= -\frac{4}{3} y_1 e^{i\theta} \beta_2^\dagger \rho^{ij} \beta_1 \\
z_3^{ij3} - 8y_3^{ij3} + i \left(z_3^{ij4} - 8y_3^{ij4} \right) &= +\frac{4}{3} y_1 e^{i\theta} \beta_3^\dagger \rho^{ij} \beta_1 \\
z_3^{ij5} - 8y_3^{ij5} + i \left(z_3^{ij6} - 8y_3^{ij6} \right) &= -\frac{4}{3} y_1 e^{i\theta} \beta_4^\dagger \rho^{ij} \beta_1
\end{aligned} \quad (4.25)$$

Using the list of matrix elements in (A.17), these equations may be solved by inspection, and we find $z_3^{12k} = z_3^{34k} = z_3^{56k} = 0$ for all k . The remaining entries may be recast most easily in terms of the matrix Z_3 defined in (2.4),

$$Z_3 = 8Y_3 + 32iy_1 e^{i\theta} \beta_1 \otimes \beta_1^t \quad (4.26)$$

This concludes the solution of the equations (3.21).

4.10 Summary of the solution

For given gauge coupling $g(x^\pi)$, the solution to the supersymmetry constraint equations (3.15) is parameterized by the undetermined coefficient, $y_1(x^\pi)$, the initial value of the supersymmetry parameter phase, $\phi(x_0^\pi)$, two real, constant, Grassmann numbers $\xi_{0\pm}$, and the three unfixed eigenvalues of the real diagonal matrix $D_3(x^\pi) = \text{diag}[1, a, b, c]$. In terms of these parameters, the supersymmetry parameter is

$$\begin{aligned} \zeta(x^\pi) &= \lambda s_+ \otimes \xi_+ + i\lambda^* s_- \otimes \xi_- \\ \xi_\pm(x^\pi) &= \xi_{0\pm} \beta_1 e^{i\theta(x^\pi)/2} \\ \theta(x^\pi) &= \theta(x_0^\pi) - 2 \int_{x_0^\pi}^{x^\pi} dx^\pi (\partial_\pi g) y_1(x^\pi) \\ (\beta_1)_a &= \delta_{1a} \end{aligned} \quad (4.27)$$

where s_+ and s_- are the two eigenvectors of $\gamma^\pi \mathcal{B}$ restricted to the $+$ chirality subspace. For every ζ that satisfies the supersymmetry equations we have two real Poincaré supersymmetries given by ζ and ζ^* . The cases with more than two supersymmetries will be discussed in the next section. The fermionic coefficients are given by

$$\begin{aligned} Y_2 &= 0 \\ Y_3 &= D_3 e^{i\theta} \end{aligned} \quad (4.28)$$

The bosonic coefficients are given by (4.20) and

$$\begin{aligned} Z_2 &= -4iy_1 (I - 4\beta_1 \otimes \beta_1^\dagger) \\ Z_3 &= 8Y_3 + 32iy_1 e^{i\theta} \beta_1 \otimes \beta_1^t \\ z_4^{ij} &= -(z_1^{ik} + z_2^{ik})(z_1^{jk} + z_2^{jk}) \end{aligned} \quad (4.29)$$

Finally the modification to the supersymmetry transformation is given by

$$\chi^i = -\left(z_1^{ij} + z_2^{ij}\right) (\rho^j)^* \gamma^\pi \mathcal{B} \zeta^* \quad (4.30)$$

Recall that we used both local and global $SU(4)$ transformations to simplify the form of Y_2 and Y_3 , and to present the solution in the simple form given above. The general solution to the supersymmetry constraint equations may finally be recovered by undoing those $SU(4)$ transformations on the above solution, namely by making local and global $SU(4)$ rotations on the above formulas.

5 Classifying interface supersymmetric theories

We now have all the tools available to give a complete classification of the allowed Poincaré supersymmetries as a function of the gauge and interface couplings. Using Lemma 1, this classification proceeds according to $r = \text{rank}(Y_3^\dagger Y_3)$, and corresponds to the following forms of Y_3 (after diagonalization by $U \in SU(4)$), following Lemma 1,

$$\begin{aligned}
r = 4 & & Y_3 &= e^{i\theta} \text{diag}[1, 1, 1, 1] \\
r = 3 & & Y_3 &= e^{i\theta} \text{diag}[1, 1, 1, c] & 0 \leq c \neq 1 \\
r = 2 & & Y_3 &= e^{i\theta} \text{diag}[1, 1, b, c] & 0 \leq b \neq 1 \\
r = 1 & & Y_3 &= e^{i\theta} \text{diag}[1, a, b, c] & 0 \leq a \neq 1
\end{aligned} \tag{5.1}$$

Within a category with fixed amount of supersymmetry (i.e. fixed r), we shall principally be interested in the theory which has *maximal internal symmetry*, since other theories with the same amount of supersymmetry but less internal symmetries may be viewed as perturbations of the former by BPS operators that further break the internal symmetry.

5.1 Solutions with extended interface supersymmetry

Extended interface supersymmetry ($r > 1$) will occur when solutions exist to the equations (3.21), for given interface couplings, for at least two linearly independent unit vectors β . By $SU(4)$ symmetry, these vectors may be chosen to be β_1 and β_2 , so that Y_3 is of the form (5.1) with $r = 2$. The cases with $r = 3, 4$ are contained in this case by setting b and possibly c equal to 1.

For a given interface theory, the interface couplings Z_2 are fixed. The simultaneous solution for β_1 and β_2 will require that the expression (4.23) for Z_2 hold in terms of these two independent vectors,

$$\begin{aligned}
Z_2 &= -4iy_1 (I - 4\beta_1 \otimes \beta_1^\dagger) \\
&= -4iy_1 (I - 4\beta_2 \otimes \beta_2^\dagger)
\end{aligned} \tag{5.2}$$

These equations can hold only provided $y_1 = 0$, so that,

$$y_1 = Y_2 = Z_2 = 0 \qquad Z_3 = 8Y_3 \tag{5.3}$$

which provides a considerable simplification of the solution, and renders the supersymmetry parameter ξ independent of x^π . One can check that all of the supersymmetry equations in (3.21) hold, except that (2') imposes the following consistency condition,

$$b = c e^{i4\theta} \tag{5.4}$$

By construction of D_3 , we have $b, c \geq 0$, so the solutions fall into three distinct classes,

(I)	$b = c = 1$	$e^{4i\theta} = 1$	$SU(2) \times SU(2)$
(II)	$b = c = 0$	θ arbitrary	$SO(2) \times SU(2)$
(III)	$b = c \neq 0, 1$	$e^{4i\theta} = 1$	$SO(2) \times SO(2)$

(5.5)

On the right side, we have listed the residual symmetry of the solution. From these considerations, it is immediately clear that the case $r = 3$ does not support any solutions, since it would have $c \neq b = 1$.

5.2 The theory with $\mathcal{N} = 4$ interface supersymmetry

Case (I) corresponds to a theory with 8 conserved Poincaré supercharges, or $\mathcal{N} = 4$ *interface supersymmetry*, and $SU(2) \times SU(2) \sim SO(4)$ R-symmetry. Setting first the $SU(4)$ rotation matrix U equal to 1, the interface Lagrangian is given by⁸

$$\begin{aligned} \mathbb{L}_I = & \frac{(\partial_\pi g)}{g^3} \text{tr} \left(\frac{i}{2} \psi^t \mathcal{C} \psi + \frac{i}{2} \psi^\dagger \mathcal{C} \psi^* - 4i \phi^2 [\phi^4, \phi^6] + \partial_\pi (\phi^2 \phi^2 + \phi^4 \phi^4 + \phi^6 \phi^6) \right) \\ & - \frac{(\partial_\pi g)^2}{g^4} 2 \text{tr} (\phi^2 \phi^2 + \phi^4 \phi^4 + \phi^6 \phi^6) \end{aligned} \quad (5.6)$$

Using (6.4), we can scale out the $(\partial_\pi g)^2$ term by defining $\tilde{\phi}^i = \phi^i / g^2$ for i even. Our conventions are as follows, $\tilde{\phi}^i$ has $i \in \{2, 4, 6\}$ and ϕ^i has $i \in \{1, 3, 5\}$. In terms of these fields the bulk Lagrangian is

$$\begin{aligned} \mathbb{L}_0 = & -\frac{1}{4g^2} \text{tr} (F^{\mu\nu} F_{\mu\nu}) - \frac{1}{2g^2} \text{tr} (D^\mu \phi^i D_\mu \phi^i) - \frac{g^2}{2} \text{tr} (D^\mu \tilde{\phi}^i D_\mu \tilde{\phi}^i) \\ & + \frac{1}{4g^2} \text{tr} ([\phi^i, \phi^j] [\phi^i, \phi^j]) + \frac{g^2}{2} \text{tr} ([\tilde{\phi}^i, \phi^j] [\tilde{\phi}^i, \phi^j]) + \frac{g^6}{4} \text{tr} ([\tilde{\phi}^i, \tilde{\phi}^j] [\tilde{\phi}^i, \tilde{\phi}^j]) \\ & - \frac{i}{2g^2} \text{tr} (\bar{\psi} \gamma^\mu D_\mu \psi) + \frac{i}{2g^2} \text{tr} (D_\mu \bar{\psi} \gamma^\mu \psi) + \frac{1}{2g^2} \text{tr} (\psi^t \mathcal{C} \rho^i [\phi^i, \psi] + \psi^\dagger \mathcal{C} (\rho^i)^* [\phi^i, \psi^*]) \\ & + \frac{1}{2} \text{tr} (\psi^t \mathcal{C} \rho^i [\tilde{\phi}^i, \psi] + \psi^\dagger \mathcal{C} (\rho^i)^* [\tilde{\phi}^i, \psi^*]) \end{aligned} \quad (5.7)$$

and the interface Lagrangian is

$$\mathbb{L}_I = \frac{(\partial_\pi g)}{g^3} \text{tr} \left(\frac{i}{2} \psi^t \mathcal{C} \psi + \frac{i}{2} \psi^\dagger \mathcal{C} \psi^* - 4ig^6 \tilde{\phi}^2 [\tilde{\phi}^4, \tilde{\phi}^6] \right) \quad (5.8)$$

⁸The cases with $\theta = \pm\pi/2, \pi$ are equivalent to $\theta = 0$, which is the choice made here.

The $SU(2) \times SU(2) \subset SU(4)$ symmetry acts as follows. The first $SU(2)$ transforms the scalar triplet (ϕ^1, ϕ^3, ϕ^5) in its 3-dimensional representation, while the second $SU(2)$ transforms the scalar triplet (ϕ^2, ϕ^4, ϕ^6) . On the fermion ψ , the group $SU(2) \times SU(2) \sim SO(4)$ acts in the (real) 4-dimensional representation of $SO(4)$. This theory clearly admits a conformal limit, as all interface couplings proportional to $(\partial_\pi g)^2$ have been eliminated. Therefore, the 8 Poincaré supercharges are supplemented by 8 more conformal supercharges to a total of 16 conformal supersymmetries.

Finally, we may restore the dependence on the $SU(4)$ rotation matrix U by including it in the form of Y_3 and $Z_3 = 8Y_3$, which gives us the general interface Lagrangian with $\mathcal{N} = 4$ interface supersymmetry,

$$\mathcal{L}_I = \frac{(\partial_\pi g)}{g^3} \text{tr} \left(\frac{i}{2} \psi^t \mathcal{C} U^* U^\dagger \psi + \frac{i}{2} \psi^\dagger \mathcal{C} U U^t \psi^* + i g^6 \text{Re tr} \left(\rho^{ijk} U U^t \right) \tilde{\phi}^i [\tilde{\phi}^j, \tilde{\phi}^k] \right) \quad (5.9)$$

where we have used (A.12) in re-expressing the last term. The space of interface theories is thus parametrized by the gauge coupling and the interface couplings

$$U \in \frac{SU(4)}{SO(4)} \quad (5.10)$$

where $SO(4)$ acts on U by right multiplication. Theories for different U are physically equivalent, although described by a different set of couplings.

5.3 The theories with $\mathcal{N} = 2$ interface supersymmetry

Cases (II) and (III) correspond to theories which have 4 conserved Poincaré supercharges, or $\mathcal{N} = 2$ *interface supersymmetry*. Case (II) has $SO(2) \times SU(2)$ R-symmetry, while (III) has a smaller internal symmetry $SO(2) \times SO(2)$. The theory corresponding to case (III) may be obtained from the theory corresponding to case (II) by adding interface operators that preserve the supersymmetry but further break the internal symmetry. We shall describe in detail only the theory with maximal internal symmetry, i.e. case (II). Since D_3 has zero eigenvalues, by Lemma 1, we may set $\theta = 0$ without loss of generality, since the action of this angle is equivalent to an $SU(4)$ rotation. Setting first the $SU(4)$ rotation matrix U to 1, the interface Lagrangian for theory (II) is given by,

$$\begin{aligned} \mathcal{L}_I = & \frac{(\partial_\pi g)}{g^3} \text{tr} \left(\frac{i}{2} \psi^t \mathcal{C} D_3^{(2)} \psi + \frac{i}{2} \psi^\dagger \mathcal{C} D_3^{(2)} \psi^* + 2i \phi^2 [\phi^3, \phi^5] - 2i \phi^2 [\phi^4, \phi^6] \right. \\ & \left. + \frac{1}{2} \partial_\pi (\phi^i \phi^i + \phi^2 \phi^2 - \phi^1 \phi^1) \right) \\ & - \frac{(\partial_\pi g)^2}{2g^4} \text{tr} \left(\phi^i \phi^i + 3 \phi^2 \phi^2 - \phi^1 \phi^1 \right) \end{aligned} \quad (5.11)$$

where $D_3^{(2)} = \text{diag}[1, 1, 0, 0]$. Using (6.4), we can scale out the $(\partial_\pi g)^2$ term by defining $\tilde{\phi}^2 = \frac{\phi^2}{g^2}$ and $\tilde{\phi}^i = \frac{\phi^i}{g}$ for $i \in \{3, \dots, 6\}$. In terms of $\tilde{\phi}^i$, the interface Lagrangian is

$$\mathbb{L}_I = \frac{(\partial_\pi g)}{g^3} \text{tr} \left(\frac{i}{2} \psi^t \mathcal{C} D_3^{(2)} \psi + \frac{i}{2} \psi^\dagger \mathcal{C} D_3^{(2)} \psi^* + g^4 2i \tilde{\phi}^2 [\tilde{\phi}^3, \tilde{\phi}^5] - g^4 2i \tilde{\phi}^2 [\tilde{\phi}^4, \tilde{\phi}^6] \right) \quad (5.12)$$

In this form, it is clear that these interface theories admit a conformal limit, so that the 4 Poincaré supersymmetries get enhanced with an extra 4 conformal supersymmetries.

Restoring the dependence of the interface couplings on the matrix U gives the general interface Lagrangian with $\mathcal{N} = 2$ interface supersymmetry,

$$\begin{aligned} \mathbb{L}_I = & \frac{(\partial_\pi g)}{g^3} \text{tr} \left(\frac{i}{2} \psi^t \mathcal{C} U^* D_3^{(2)} U^\dagger \psi + \frac{i}{2} \psi^\dagger \mathcal{C} U D_3^{(2)} U^t \psi^* \right. \\ & \left. + i g^4 \text{Re tr} \left(\rho^{ijk} U D_3^{(2)} U^t \right) \tilde{\phi}^i [\tilde{\phi}^j, \tilde{\phi}^k] \right) \end{aligned} \quad (5.13)$$

The space of $\mathcal{N} = 2$ interface theories is thus parametrized by the interface couplings

$$U \in \frac{SU(4)}{SO(2) \times SU(2)} \quad (5.14)$$

where $SO(2) \times SU(2)$ acts on U by right multiplication, and leaves $D_3^{(2)}$ invariant. Theories for different U are again physically equivalent, although described by different interface couplings.

5.4 The theory with $\mathcal{N} = 1$ interface supersymmetry

Finally, the case $r = 1$ in (5.1) corresponds to a family of theories with 2 conserved Poincaré supercharges, or $\mathcal{N} = 1$ *interface supersymmetry*. Its maximal internal symmetry is $SU(3)$ attained in the case where $a = b = c = 0$, while no internal symmetry remains when a, b, c are all distinct. We shall describe in detail only the theory with maximal internal symmetry $SU(3)$ and $a = b = c = 0$. Since D_3 has zero eigenvalues, by Lemma 1, we may set $\theta = 0$ without loss of generality. Using the results of (6.6), we take $y_1 = 0$, so that we may scale out the $(\partial_\pi g)^2$ term like before. Setting first the $SU(4)$ rotation matrix to 1, the interface Lagrangian for this theory is given by,

$$\begin{aligned} \mathbb{L}_I = & \frac{(\partial_\pi g)}{g^3} \text{tr} \left(\frac{i}{2} \psi^t \mathcal{C} D_3^{(1)} \psi + \frac{i}{2} \psi^\dagger \mathcal{C} D_3^{(1)} \psi^* \right) \\ & + \frac{(\partial_\pi g)}{g^3} \text{tr} \left(i \phi^1 [\phi^3, \phi^6] + i \phi^1 [\phi^4, \phi^5] + i \phi^2 [\phi^3, \phi^5] \right. \\ & \left. - i \phi^2 [\phi^4, \phi^6] + \frac{1}{2} \partial_\pi (\phi^i \phi^i) \right) - \frac{(\partial_\pi g)^2}{2g^4} \text{tr} (\phi^i \phi^i) \end{aligned} \quad (5.15)$$

where $D_3^{(1)} = \text{diag}[1, 0, 0, 0]$. We can scale out the $(\partial_\pi g)^2$ term by defining $\tilde{\phi}^i = \phi^i/g$. In terms of $\tilde{\phi}^i$ the interface Lagrangian is

$$\begin{aligned} \mathbb{L}_I = & \frac{(\partial_\pi g)}{g^3} \text{tr} \left(\frac{i}{2} \psi^t \mathcal{C} D_3^{(1)} \psi + \frac{i}{2} \psi^\dagger \mathcal{C} D_3^{(1)} \psi^* \right) \\ & + (\partial_\pi g) \text{tr} \left(i \tilde{\phi}^1 [\tilde{\phi}^3, \tilde{\phi}^6] + i \tilde{\phi}^1 [\tilde{\phi}^4, \tilde{\phi}^5] + i \tilde{\phi}^2 [\tilde{\phi}^3, \tilde{\phi}^5] - i \tilde{\phi}^2 [\tilde{\phi}^4, \tilde{\phi}^6] \right) \end{aligned} \quad (5.16)$$

In this formulation, it is clear that the theory admits a conformal limit, so that the 2 Poincaré supersymmetries get enhanced by an extra 2 conformal supersymmetries. Notice, however, that the rescaling powers of the gauge coupling g are now different from those needed for the theories with $\mathcal{N} = 4$ and $\mathcal{N} = 2$ interface supersymmetry.

In terms of the complex fields

$$\begin{aligned} \Phi_1 &= \tilde{\phi}^1 + i \tilde{\phi}^2 & \Phi_1^* &= \tilde{\phi}^1 - i \tilde{\phi}^2 \\ \Phi_2 &= \tilde{\phi}^3 + i \tilde{\phi}^4 & \Phi_2^* &= \tilde{\phi}^3 - i \tilde{\phi}^4 \\ \Phi_3 &= \tilde{\phi}^5 + i \tilde{\phi}^6 & \Phi_3^* &= \tilde{\phi}^5 - i \tilde{\phi}^6 \end{aligned} \quad (5.17)$$

the Lagrangian may be recast in a simplified form,

$$\mathbb{L}_I = \frac{(\partial_\pi g)}{g^3} \text{tr} \left(\frac{i}{2} \psi^t \mathcal{C} D_3^{(1)} \psi + \frac{i}{2} \psi^\dagger \mathcal{C} D_3^{(1)} \psi^* + \frac{g^3}{2} \Phi_1 [\Phi_2, \Phi_3] - \frac{g^3}{2} \Phi_1^* [\Phi_2^*, \Phi_3^*] \right) \quad (5.18)$$

In this form, the $SU(3)$ symmetry is manifest on the scalar fields as well as on the fermions, and rotates the fields Φ_1, Φ_2 , and Φ_3 into one another.

Restoring the dependence of the interface couplings on the matrix U gives the general interface Lagrangian with $\mathcal{N} = 1$ interface supersymmetry,

$$\begin{aligned} \mathbb{L}_I = & \frac{(\partial_\pi g)}{g^3} \text{tr} \left(\frac{i}{2} \psi^t \mathcal{C} U^* D_3^{(1)} U^\dagger \psi + \frac{i}{2} \psi^\dagger \mathcal{C} U D_3^{(1)} U^t \psi^* \right. \\ & \left. + i g^3 \text{Re tr} \left(\rho^{ijk} U D_3^{(1)} U^t \right) \tilde{\phi}^i [\tilde{\phi}^j, \tilde{\phi}^k] \right) \end{aligned} \quad (5.19)$$

The space of physically equivalent $\mathcal{N} = 1$ interface theories is thus parameterized by the interface couplings

$$U \in \frac{SU(4)}{SU(3)} \quad (5.20)$$

where $SU(3)$ acts on U by right multiplication, and leaves $D_3^{(1)}$ invariant. Of this 7-dimensional space of interface couplings, 6 may be absorbed by taking suitably modified

linear combinations for the definition of the fields Φ_1 , Φ_2 , and Φ_3 in (5.17). The remaining diagonal generator of $SU(4)$ leaves the following angle,

$$\begin{aligned} \mathbb{L}_I = & \frac{(\partial_\pi g)}{g^3} \text{tr} \left(\frac{i}{2} e^{+3i\theta} \psi^t \mathcal{C} D_3^{(1)} \psi + \frac{i}{2} e^{-3i\theta} \psi^\dagger \mathcal{C} D_3^{(1)} \psi^* \right. \\ & \left. + \frac{g^3}{2} e^{+3i\theta} \Phi_1 [\Phi_2, \Phi_3] - \frac{g^3}{2} e^{-3i\theta} \Phi_1^* [\Phi_2^*, \Phi_3^*] \right) \end{aligned} \quad (5.21)$$

In [11] this Lagrangian was constructed using the language of $\mathcal{N} = 1$ theories. It was necessary to use different scalings for the chiral and vector multiplets. We now see that the origin of this lies in eliminating the z_4 term proportional to $(\partial_\pi g)^2$ in order to guarantee the existence of a conformal limit. Alternatively, one may keep the same scaling for the chiral and vector multiplets but obscure conformal invariance.

5.5 Summary of symmetry considerations

In the interface theory, the $SU(4)$ transformations of the original $\mathcal{N} = 4$ super Yang-Mills naturally fall into three distinct classes.

Transformations in the first class form the subgroup G of $SU(4)$ which leaves D_3 of (4.13) and the supersymmetry, or $\xi_\pm$, invariant.⁹ It follows that G will leave all other interface couplings invariant, thus yielding an *internal* symmetry of the full interface theory. To see this note that invariance of Z_2 , Z_3 , Y_2 and Y_3 follows from (4.28), (4.29) and the transformation rules (4.4); z_1^{ij} is completely determined by Y_3 and $\xi_\pm$ which are invariant; invariance of z_4^{ij} follows from invariance of z_1^{ij} and z_2^{ij} . These groups are $G = SU(3)$, $G = SU(2)$ and the identity group $G = \{I\}$ for the theories with 2, 4, and 8 Poincaré supercharges respectively.

Transformations in the second class form a subgroup of $SU(4)$ which leaves the interface couplings invariant, but transforms the supersymmetries into one another, thus yielding the R-symmetry of the full interface theory. The R-symmetry group turns out to be $SO(n)$, where n is the number of linearly independent ξ that satisfy the supersymmetry constraint equations (3.21). The space-time symmetries may then be conveniently packaged in the supergroup $OSP(2, 2|n)$.

Transformations in the third class are dualities generated by $U \in SU(4)$ which do not lie in G or $SO(n)$. These transformations change the interface couplings, and thus

⁹In the case of more than one supersymmetry, we take G to leave invariant all of the supersymmetries as well as D_3 .

provide a map between interface theories related by $SU(4)$ rotations. Since the supersymmetry constraint equations are covariant under $SU(4)$, these different interface theories will have the same amount of supersymmetry. Such transformations are parameterized by the coset

$$\frac{SU(4)}{G \times SO(n)} \quad (5.22)$$

and relate different, but physically equivalent theories. This is analogous to Montonen-Olive duality, which states that two different theories related by $SL(2, \mathbf{Z})$ transformations are physically equivalent. A special role is played by transformations $U \in SU(4)/(G \times SO(n))$ which do not change the embedding of $G \times SO(n)$. These transformations form the subgroup H of $SU(4)$ which commutes with $G \times SO(n)$.

For the AdS dual, not all of the $SU(4)$ transformations can be implemented in a useful way. First, one can choose an Ansatz that is invariant under $G \times SO(n)$. Once this is done, the embedding of $G \times SO(n)$ is fixed and the only useful transformations of $SU(4)$ that can be implemented on the $G \times SO(n)$ -invariant Ansatz are those in H . Since we did not include $\text{tr}(F\tilde{F})$ or L_{CS} , given in (2.5), in our analysis, our theory cannot exhibit Montonen-Olive duality. We expect that upon including these terms, we should recover Montonen-Olive duality, which degenerates to $SL(2, \mathbf{R})$ in the supergravity limit. This gives the following symmetry Ansatz for the AdS dual

$$SO(2, 3) \times SO(n) \times G \times H \times SL(2, \mathbf{R}) \quad (5.23)$$

Based on these symmetry considerations for the Ansatz of the AdS dual, an explicit ten dimensional supergravity solution is found in [2], with $G = SU(3)$, and the solution generating symmetries $U(1)_\beta \times SL(2, \mathbf{R})$.

6 Conformal symmetry

We now analyze under what conditions conformal symmetry may be recovered. First note that the gauge coupling needs to be invariant under a scale transformation. One way to achieve this is to take the gauge coupling to be a step function. However, the z_4 term in the interface Lagrangian (2.3) is proportional to $(\partial_\pi g)^2$ and does not have a well-defined limit. In some cases it is possible to eliminate z_4 by rescaling the scalar fields. An example of this was already shown in [11]. We examine the conditions necessary for such a rescaling to work in general. We define rescaled scalar fields $\tilde{\phi}^i$ by $\phi^i = g^{n_i} \tilde{\phi}^i$, where n_i is the real scaling exponent which is to be solved for. The relevant part of the Lagrangian is

$$\mathcal{L}_1 = -\frac{1}{2g^2} \text{tr} \left(D_\mu \phi^i D^\mu \phi^i \right) + \frac{(\partial_\pi g)}{2g^3} z_1^{ij} \partial_\pi \text{tr} \left(\phi^i \phi^j \right) + \frac{(\partial_\pi g)^2}{2g^4} z_4^{ij} \text{tr} \left(\phi^i \phi^j \right) \quad (6.1)$$

In terms of rescaled fields the Lagrangian becomes

$$\begin{aligned} \mathcal{L}_1 = & -\frac{g^{2n_i}}{2g^2} \text{tr} \left(D_\mu \tilde{\phi}^i D^\mu \tilde{\phi}^i \right) - \frac{g^{n_i+n_j}}{2g^3} \left(n_i \delta_{ij} - z_1^{ij} \right) (\partial_\pi g) \partial_\pi \text{tr} \left(\tilde{\phi}^i \tilde{\phi}^j \right) \\ & - \frac{g^{n_i+n_j}}{2g^4} \left(n_i^2 \delta_{ij} - (n_i + n_j) z_1^{ij} - z_4^{ij} \right) (\partial_\pi g)^2 \text{tr} \left(\tilde{\phi}^i \tilde{\phi}^j \right) \end{aligned} \quad (6.2)$$

The conformal limit will exist, provided there exists a rescaling $\phi^i = g^{n_i} \tilde{\phi}^i$ such that the term proportional to $(\partial_\pi g)^2$ vanishes; this can be achieved when

$$n_i^2 \delta_{ij} - (n_i + n_j) z_1^{ij} - z_4^{ij} = 0 \quad (6.3)$$

For $i = j$, we obtain 6 equations which determine the exponents n_i , as follows,

$$\begin{aligned} 0 &= -n_i^2 + 2n_i z_1^{ii} + z_4^{ii} \\ n_i &= z_1^{ii} \pm \sqrt{(z_1^{ii})^2 + z_4^{ii}} \end{aligned} \quad (6.4)$$

Using the explicit form for z_4^{ij} , obtained from the supersymmetry condition (3.22), we deduce a simple expression for the argument of the square root in the formula for n_i ,

$$(z_1^{ii})^2 + z_4^{ii} = - \sum_{k \neq i} \left(z_1^{ik} + z_2^{ik} \right)^2 \quad (6.5)$$

Reality of n_i requires these arguments to be non-negative, which can hold only when,

$$z_1^{ik} = z_2^{ik} = 0 \quad i \neq k \quad (6.6)$$

These results imply that $y_1 = 0$ along with either $\sin(2\theta) = 0$ and/or $a = b = c = 0$.

7 Comparing with duals to brane configurations

As was remarked in the Introduction, no compelling brane configuration candidate is available yet which naturally reduces to the non-supersymmetric Janus solution in the near-horizon limit. It may be hoped that, with the additional restrictions provided by supersymmetry, a comparison between candidate brane configurations and interface dual Yang-Mills theories will be considerably facilitated. The more supersymmetry, the more restrictive the comparison will be.

The interface Yang-Mills theory with maximal supersymmetry has $\mathcal{N} = 4$ interface supersymmetry, or 8 real Poincaré supercharges, and $SU(2) \times SU(2)$ internal symmetry. Remarkably, there exists a brane configuration which has precisely those symmetries, and yet does not produce our $\mathcal{N} = 4$ interface theory in the near-horizon limit. Indeed, a configuration with intersecting D3 and D5 branes has 8 real Poincaré supercharges, which is $1/4$ of the maximal 32 supersymmetries of the theory. This configuration also has $SU(2) \times SU(2) \sim SO(4)$ internal symmetry. In the near-horizon limit, the Poincaré symmetry gets enhanced to conformal $SO(2, 3)$, and the 8 Poincaré supercharges are supplemented with 8 special conformal supercharges, just as is the case in the conformal limit of the $\mathcal{N} = 4$ interface supersymmetry theory. Thus, the symmetries of the D3-D5 brane system and of our $\mathcal{N} = 4$ interface theory match precisely.

And yet, the supersymmetry multiplet content is different, because the D3-D5 system contains open $3 - 5$ strings stretching from the D3 to the D5 brane. Therefore, the CFT dual to the D3-D5 system is expected to include gauge degrees of freedom localized on the interface. In [22], a $2+1$ -dimensional hypermultiplet, localized on the brane, in the fundamental representation of the gauge group $SU(N)$ was proposed for such defect degrees of freedom. Ample evidence was presented there that this scenario is viable, at least in the approximation where the D5 brane is treated as a “probe”, and the back reaction of the D5 on the system of D3 branes is ignored [21, 23, 24, 25]. In our $\mathcal{N} = 4$ interface theory, however, such extra degrees of freedom are absent. If our $\mathcal{N} = 4$ interface theory is to have any relation with the D3-D5 brane system, some mechanism must exist that decouples the $3 - 5$ strings from the CFT dual theory.¹⁰ We hope to investigate these questions in the future.

¹⁰An example for such a mechanism in a different system was discovered in [26].

Acknowledgments

It is a pleasure to acknowledge helpful conversations with Iosif Bena, Per Kraus and Norisuke Sakai. This work was supported in part by National Science Foundation (NSF) grant PHY-04-56200. ED is grateful to the Kavli Institute for Theoretical Physics (KITP) for their hospitality and support under NSF grant PHY-99-07949. MG is grateful to the Harvard Particle Theory group for hospitality while this work was being completed.

A Algebra of $SU(4)$ Clebsch-Gordan coefficients

The canonical map between $SU(4)$ and $SO(6)$ is realized in terms of the Clifford algebra of $SO(6)$, given by the $\gamma_{(6)}^i$ -matrices, which obey

$$\{\gamma_{(6)}^i, \gamma_{(6)}^j\} = 2\delta^{ij}I_8 \quad i, j = 1, \dots, 6 \quad (\text{A.1})$$

The chirality matrix is defined by $\gamma_{(6)} \equiv i\gamma_{(6)}^1\gamma_{(6)}^2\cdots\gamma_{(6)}^6$ and satisfies $(\gamma_{(6)})^2 = I_8$. The charge conjugation matrix $\mathcal{C}_{(6)}$ is defined by $\mathcal{C}_{(6)}\gamma_{(6)}^i\mathcal{C}_{(6)}^{-1} = -(\gamma_{(6)}^i)^t$, and satisfies the relations $\mathcal{C}_{(6)}^t = \mathcal{C}_{(6)}^{-1} = \mathcal{C}_{(6)}^* = \mathcal{C}_{(6)}$. We choose the chiral basis of γ -matrices, where,

$$\gamma_{(6)} = \begin{pmatrix} I_4 & 0 \\ 0 & -I_4 \end{pmatrix} \quad (\text{A.2})$$

and $\mathcal{C}_{(6)}\gamma_{(6)} = -\gamma_{(6)}\mathcal{C}_{(6)}$. The chiral restriction of the Dirac matrices then defines the matrices ρ^i as follows,

$$\begin{aligned} \mathcal{C}_{(6)}\gamma_{(6)}^i &= \begin{pmatrix} \rho^i & 0 \\ 0 & -(\rho^i)^* \end{pmatrix} \\ \gamma_{(6)}^i\mathcal{C}_{(6)} &= \begin{pmatrix} -(\rho^i)^* & 0 \\ 0 & \rho^i \end{pmatrix} \end{aligned} \quad (\text{A.3})$$

We shall also need the chiral restriction of Clifford algebra generators of higher rank $SO(6)$ -tensors of the form $\gamma_{(6)}^{i_1\cdots i_p} \equiv \gamma_{(6)}^{[i_1}\cdots\gamma_{(6)}^{i_p]}$. They are given as follows,

$$\begin{aligned} \gamma_{(6)}^{ij} &= \begin{pmatrix} \rho^{ij} & 0 \\ 0 & (\rho^{ij})^* \end{pmatrix} & \rho^{ij} &\equiv -(\rho^{[i})^*\rho^{j]} \\ \mathcal{C}_{(6)}\gamma_{(6)}^{ijk} &= \begin{pmatrix} \rho^{ijk} & 0 \\ 0 & -(\rho^{ijk})^* \end{pmatrix} & \rho^{ijk} &\equiv -\rho^{[i}(\rho^j)^*\rho^{k]} \\ \gamma_{(6)}^{ijk}\mathcal{C}_{(6)} &= \begin{pmatrix} -(\rho^{ijk})^* & 0 \\ 0 & \rho^{ijk} \end{pmatrix} \end{aligned} \quad (\text{A.4})$$

Square brackets $[]$ imposes anti-symmetrization on whichever set of indices occurs inside. In view of the definition of charge conjugation, the matrices ρ^i are anti-symmetric, which implies the following properties,

$$\begin{aligned} (\rho^i)^t &= -\rho^i & \mathbf{4} \otimes \mathbf{4} &\rightarrow \mathbf{6} \\ (\rho^{ij})^\dagger &= -\rho^{ij} & \mathbf{4} \otimes \bar{\mathbf{4}} &\rightarrow \mathbf{15} \\ (\rho^{ijk})^t &= +\rho^{ijk} & \mathbf{4} \otimes \mathbf{4} &\rightarrow \mathbf{10} \end{aligned} \quad (\text{A.5})$$

These matrices provide the Clebsch-Gordan coefficients for the $SU(4)$ tensor product decompositions involving the $\mathbf{4}$ and $\bar{\mathbf{4}}$ representations, listed above in the right column.

A.1 The algebra of the ρ -matrices

The standard Clifford algebra readily provides relations between the products of ρ -matrices in various combinations. The following Clifford relations,

$$\begin{aligned}
\{\gamma^i, \gamma^j\} &= 2\delta^{ij}I \\
[\gamma^{ij}, \gamma^k] &= 2\delta^{jk}\gamma^i - 2\delta^{ik}\gamma^j \\
[\gamma^{ij}, \gamma^{kl}] &= 2\delta^{jk}\gamma^{il} + 2\delta^{jl}\gamma^{ki} - (i \leftrightarrow j) \\
[\gamma^{ij}, \gamma^{klm}] &= 2\delta^{jk}\gamma^{ilm} + 2\delta^{jl}\gamma^{kim} + 2\delta^{jm}\gamma^{kli} - (i \leftrightarrow j) \\
\{\gamma^i, \gamma^{klm}\} &= 2\delta^{ik}\gamma^{lm} + 2\delta^{il}\gamma^{mk} + 2\delta^{im}\gamma^{kl}
\end{aligned} \tag{A.6}$$

translate to the following equations for the ρ -matrices and their complex conjugates,

$$(\rho^i)^*\rho^j + (\rho^j)^*\rho^i = -2\delta^{ij}I \tag{A.7}$$

$$(\rho^{ij})^*\rho^k - \rho^k\rho^{ij} = 2\delta^{jk}\rho^i - 2\delta^{ik}\rho^j \tag{A.8}$$

$$[\rho^{ij}, \rho^{kl}] = 2\delta^{jk}\rho^{il} + 2\delta^{jl}\rho^{ki} - (i \leftrightarrow j) \tag{A.9}$$

$$(\rho^{ij})^*\rho^{klm} - \rho^{klm}\rho^{ij} = 2\delta^{jk}\rho^{ilm} + 2\delta^{jl}\rho^{kim} + 2\delta^{jm}\rho^{kii} - (i \leftrightarrow j) \tag{A.10}$$

$$(\rho^i)^*\rho^{klm} + (\rho^{klm})^*\rho^i = -2\delta^{ik}\rho^{lm} - 2\delta^{il}\rho^{mk} - 2\delta^{im}\rho^{kl} \tag{A.11}$$

These relations provide the Clebsch-Gordan coefficients for tensor products involving the **6**, **15**, **10**, and $\overline{\mathbf{10}}$ representations.

The following relations will also be useful,

$$\begin{aligned}
\text{tr}(\rho^i(\rho^{i'})^*) &= -4\delta^{ii'} \\
\text{tr}(\rho^{ijk}(\rho^{i'j'k'})^*) &= +4\delta^{i[i'}\delta^{jj'}\delta^{kk']}
\end{aligned} \tag{A.12}$$

where the bracket [] indicates anti-symmetrization of the primed indices only.

A.2 Explicit form in standard basis

The standard basis of $\gamma_{(6)}^i$ matrices yields the following convenient basis of ρ^i matrices,

$$\begin{aligned}
\rho^1 &= \sigma_3 \otimes \sigma_2 & \rho^2 &= -iI \otimes \sigma_2 \\
\rho^3 &= -\sigma_2 \otimes I & \rho^4 &= i\sigma_2 \otimes \sigma_3 \\
\rho^5 &= \sigma_1 \otimes \sigma_2 & \rho^6 &= -i\sigma_2 \otimes \sigma_1
\end{aligned} \tag{A.13}$$

In this basis, we have simple complex conjugation relations, (in addition to the general relations of (A.5)),

$$\begin{aligned}
(\rho^i)^* &= (-1)^i \rho^i \\
(\rho^{ij})^* &= (-1)^{i+j} \rho^{ij} & \rho^{ij} &= -(-1)^i \rho^i \rho^j & j &\neq i \\
(\rho^{ijk})^* &= (-1)^{i+j+k} \rho^{ijk} & \rho^{ijk} &= -(-1)^j \rho^i \rho^j \rho^k & k &\neq i, j
\end{aligned} \tag{A.14}$$

while the square of each ρ^i matrix is given by $(\rho^i)^2 = -(-1)^i$. These formulas facilitate the evaluation, in this basis, the matrices ρ^{ij} ,

$$\begin{aligned}
\rho^{12} &= -i\sigma_3 \otimes I & \rho^{23} &= -i\sigma_2 \otimes \sigma_2 & \rho^{35} &= i\sigma_3 \otimes \sigma_2 \\
\rho^{13} &= i\sigma_1 \otimes \sigma_2 & \rho^{24} &= -i\sigma_2 \otimes \sigma_1 & \rho^{36} &= iI \otimes \sigma_1 \\
\rho^{14} &= i\sigma_1 \otimes \sigma_1 & \rho^{25} &= i\sigma_1 \otimes I & \rho^{45} &= i\sigma_3 \otimes \sigma_1 \\
\rho^{15} &= i\sigma_2 \otimes I & \rho^{26} &= -i\sigma_2 \otimes \sigma_3 & \rho^{46} &= -iI \otimes \sigma_2 \\
\rho^{16} &= i\sigma_1 \otimes \sigma_3 & \rho^{34} &= -iI \otimes \sigma_3 & \rho^{56} &= -i\sigma_3 \otimes \sigma_3
\end{aligned} \tag{A.15}$$

and the matrices ρ^{ijk} ,

$$\begin{aligned}
\rho^{123} &= +i\rho^{456} = -\sigma_1 \otimes I & \rho^{135} &= -i\rho^{246} = iI \otimes I \\
\rho^{124} &= -i\rho^{356} = i\sigma_1 \otimes \sigma_3 & \rho^{136} &= +i\rho^{245} = \sigma_3 \otimes \sigma_3 \\
\rho^{125} &= +i\rho^{346} = -\sigma_2 \otimes \sigma_2 & \rho^{145} &= +i\rho^{236} = I \otimes \sigma_3 \\
\rho^{126} &= -i\rho^{345} = -i\sigma_1 \otimes \sigma_1 & \rho^{146} &= -i\rho^{235} = -i\sigma_3 \otimes I \\
\rho^{134} &= +i\rho^{256} = \sigma_3 \otimes \sigma_1 & \rho^{156} &= -i\rho^{234} = I \otimes \sigma_1
\end{aligned} \tag{A.16}$$

By inspection, it is now manifest that ρ^i , ρ^{ij} and ρ^{ijk} provide a basis respectively for complex antisymmetric, antihermitian, and complex symmetric traceless 4×4 matrices, confirming the Clebsch-Gordan maps of (A.5).

Finally, the following matrix elements will be useful when solving (3'),

$$\begin{aligned}
\rho^{12}\beta_1 &= -i\beta_1 & \rho^{13}\beta_1 &= -\beta_4 & \rho^{14}\beta_1 &= i\beta_4 \\
\rho^{15}\beta_1 &= -\beta_3 & \rho^{16}\beta_1 &= i\beta_3 & \rho^{23}\beta_1 &= i\beta_4 \\
\rho^{24}\beta_1 &= +\beta_4 & \rho^{25}\beta_1 &= i\beta_3 & \rho^{26}\beta_1 &= +\beta_3 \\
\rho^{34}\beta_1 &= -i\beta_1 & \rho^{35}\beta_1 &= -\beta_2 & \rho^{36}\beta_1 &= i\beta_2 \\
\rho^{45}\beta_1 &= i\beta_2 & \rho^{46}\beta_1 &= +\beta_2 & \rho^{56}\beta_1 &= -i\beta_1
\end{aligned} \tag{A.17}$$

References

- [1] D. Bak, M. Gutperle and S. Hirano, “A dilatonic deformation of $\text{AdS}(5)$ and its field theory dual,” *JHEP* **0305**, 072 (2003) [arXiv:hep-th/0304129].
- [2] E. D’Hoker, J. Estes and M. Gutperle, “Ten-dimensional supersymmetric Janus solutions”, UCLA/06/TEP/02 preprint (2006), [hep-th/0603012]
- [3] D. Z. Freedman, C. Nunez, M. Schnabl and K. Skenderis, “Fake supergravity and domain wall stability,” *Phys. Rev. D* **69**, 104027 (2004) [arXiv:hep-th/0312055].
- [4] A. Celi, A. Ceresole, G. Dall’Agata, A. Van Proeyen and M. Zagermann, “On the fakeness of fake supergravity,” *Phys. Rev. D* **71**, 045009 (2005) [arXiv:hep-th/0410126].
- [5] J. M. Maldacena, “The large N limit of superconformal field theories and supergravity,” *Adv. Theor. Math. Phys.* **2**, 231 (1998) [*Int. J. Theor. Phys.* **38**, 1113 (1999)] [arXiv:hep-th/9711200].
- [6] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, “Gauge theory correlators from non-critical string theory,” *Phys. Lett. B* **428**, 105 (1998) [arXiv:hep-th/9802109].
- [7] E. Witten, “Anti-de Sitter space and holography,” *Adv. Theor. Math. Phys.* **2**, 253 (1998) [arXiv:hep-th/9802150].
- [8] E. D’Hoker and D. Z. Freedman, “Supersymmetric gauge theories and the AdS/CFT correspondence,” arXiv:hep-th/0201253.
- [9] O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri and Y. Oz, “Large N field theories, string theory and gravity,” *Phys. Rept.* **323** (2000) 183 [arXiv:hep-th/9905111].
- [10] I. Papadimitriou and K. Skenderis, “Correlation functions in holographic RG flows,” *JHEP* **0410**, 075 (2004) [arXiv:hep-th/0407071].
- [11] A. B. Clark, D. Z. Freedman, A. Karch and M. Schnabl, “The dual of Janus $((\mathfrak{g}_1)_+ \mathfrak{g}_2)_\pm$ an interface CFT,” *Phys. Rev. D* **71**, 066003 (2005) [arXiv:hep-th/0407073].
- [12] J. L. Cardy, “Conformal Invariance And Surface Critical Behavior,” *Nucl. Phys. B* **240** (1984) 514.
- [13] J. L. Cardy and D. C. Lewellen, “Bulk and boundary operators in conformal field theory,” *Phys. Lett. B* **259**, 274 (1991).
- [14] D. M. McAvity and H. Osborn, “Conformal field theories near a boundary in general dimensions,” *Nucl. Phys. B* **455** (1995) 522 [arXiv:cond-mat/9505127].
- [15]

- [15] M. Eto, Y. Isozumi, M. Nitta, K. Ohashi, K. Ohta and N. Sakai, “D-brane configurations for domain walls and their webs,” AIP Conf. Proc. **805**, 354 (2005) [arXiv:hep-th/0509127].
- [16] M. Eto, Y. Isozumi, M. Nitta, K. Ohashi and N. Sakai, “Webs of walls,” Phys. Rev. D **72**, 085004 (2005) [arXiv:hep-th/0506135].
- [17] H. Oda, K. Ito, M. Naganuma and N. Sakai, “An exact solution of BPS domain wall junction,” Phys. Lett. B **471**, 140 (1999) [arXiv:hep-th/9910095].
- [18] A. Clark and A. Karch, “Super Janus,” JHEP **0510** (2005) 094 [arXiv:hep-th/0506265].
- [19] L. Brink, J. H. Schwarz and J. Scherk, “Supersymmetric Yang-Mills Theories,” Nucl. Phys. B **121**, 77 (1977).
- [20] F. Gliozzi, J. Scherk and D. I. Olive, “Supersymmetry, Supergravity Theories And The Dual Spinor Model,” Nucl. Phys. B **122**, 253 (1977).
- [21] A. Karch and L. Randall, “Open and closed string interpretation of SUSY CFT’s on branes with boundaries,” JHEP **0106** (2001) 063 [arXiv:hep-th/0105132].
- [22] O. DeWolfe, D. Z. Freedman and H. Ooguri, “Holography and defect conformal field theories,” Phys. Rev. D **66**, 025009 (2002) [arXiv:hep-th/0111135].
- [23] O. Aharony, O. DeWolfe, D. Z. Freedman and A. Karch, “Defect conformal field theory and locally localized gravity,” JHEP **0307** (2003) 030 [arXiv:hep-th/0303249].
- [24] J. Erdmenger, Z. Guralnik and I. Kirsch, “Four-dimensional superconformal theories with interacting boundaries or defects,” Phys. Rev. D **66** (2002) 025020 [arXiv:hep-th/0203020].
- [25] S. Yamaguchi, “AdS branes corresponding to superconformal defects,” JHEP **0306** (2003) 002 [arXiv:hep-th/0305007].
- [26] N. Itzhaki, D. Kutasov and N. Seiberg, “I-brane dynamics,” JHEP **0601** (2006) 119 [arXiv:hep-th/0508025].